

Bosonization of ZF Algebras: Direction Toward Deformed Virasoro Algebra

Sergei Lukyanov[†] and Yaroslav Pugai

Department of Physics and Astronomy
Rutgers University, Piscataway, NJ 08855-049, USA

and

L.D. Landau Institute for Theoretical Physics
Chernogolovka, 142432, Russia

Abstract

These lectures were prepared to be presented at A.A. Belavin seminar on CFT at Landau Institute for Theoretical Physics. We review bosonization of CFT and show how it can be applied to the studying of representations of Zamolodchikov-Faddeev (ZF) algebras. In the bosonic construction we obtain explicit realization of chiral vertex operators interpolating between irreducible representations of the deformed Virasoro algebra. The commutation relations of these operators are determined by the elliptic matrix of IRF type and their matrix elements are given in the form of the contour integrals of some meromorphic functions.

May, 94

[†] Current address after September 1, 1994: Newman Laboratory, Cornell University, Ithaca, NY 14853-5001

1. Introduction.

The development of CFT was initiated in the fundamental work [1] by Belavin, Polyakov and Zamolodchikov where the system of axioms describing CFT was proposed. The main idea of BPZ is that fields in CFT are classified by irreducible representations of the Virasoro algebra. From mathematical point of view, the studying of the CFT is equivalent to the description of representations of the Virasoro algebra and deriving the matrix elements of vertex operators interpolating between different irreducible representations of the Virasoro algebra $\mathcal{L}_\lambda$ specified by highest weight Δ_λ . Among vertex operators there is a set of basic ones, called primary operators

$$\Phi_\Delta^{\lambda\ \mu}(\zeta) : \mathcal{L}_\mu \rightarrow \mathcal{L}_\lambda \otimes \mathbf{C}(\zeta) \zeta^{\Delta_\lambda - \Delta_\mu} .$$

Matrix elements of these operators ("conformal blocks") are multivalued analytical functions. Knowing conformal blocks one can reconstruct physical correlation functions which satisfy the requirement of locality. The analytical properties of conformal blocks [2] are dictated by the commutation relations in the algebra of chiral vertex operators $\Phi_\Delta^{\lambda\ \mu}(\zeta)$:

$$\Phi_{\Delta_1}^{\lambda_3\ \lambda_4}(\zeta_1) \Phi_{\Delta_2}^{\lambda_4\ \lambda_1}(\zeta_2)|_{\mathcal{L}_{\lambda_1}} = \sum_{\lambda_2} \mathbf{W}_{\Delta_1 \Delta_2} \begin{bmatrix} \lambda_3 & \lambda_2 \\ \lambda_4 & \lambda_1 \end{bmatrix} \Phi_{\Delta_2}^{\lambda_3\ \lambda_2}(\zeta_2) \Phi_{\Delta_1}^{\lambda_2\ \lambda_1}(\zeta_1)|_{\mathcal{L}_{\lambda_1}} . \quad (1.1)$$

This quadratic algebra turns out to be associative, which follows from the fact that matrix $\mathbf{W}_{\Delta_1 \Delta_2}$ satisfies to the Yang-Baxter equation (YBE) in the IRF form [3].

It is important that CFT can be alternatively considered as a representation theory of this algebra [4], [5] which will be called further as Zamolodchikov-Faddeev (ZF) algebra of IRF type. In this way Virasoro algebra can be considered as an algebra of transformations preserving the commutation relations. Under the appropriate choice of the set of irreducible representations of Virasoro algebra $\mathcal{L}_\lambda$ the representations of the chiral vertex algebra (1.1) possess the realization in the direct sum $\oplus_\lambda \mathcal{L}_\lambda$. ZF algebra of IRF type is deeply connected with another associative quadratic algebra

$$Z_a(\zeta_1) Z_b(\zeta_2) = \mathbf{R}_{ab}^{cd} Z_d(\zeta_2) Z_c(\zeta_1) , \quad (1.2)$$

where matrix $\mathbf{R}_{cd}^{ab}$ is a solution of the Yang-Baxter equation. We call (1.2) by Zamolodchikov-Faddeev algebra [6], [7] of vertex type. It can be realized in the extension of the space $\oplus_\lambda \mathcal{L}_\lambda$ by taking irreducible representations of Virasoro algebra with proper multiplicities:

$$\pi_Z = \oplus_\lambda \mathcal{L}_\lambda \otimes \mathcal{V}_\lambda , \quad (1.3)$$

where $\mathcal{V}_\lambda$ are some finite-dimensional vector spaces [8].

Thus, CFT can be described basing on the notions of two algebras which have different forms and require, at the first glance, different approaches of the investigation. The first one is infinite-dimensional Virasoro algebra, while the second is associative quadratic algebra which is determined by some finite-dimensional matrices. However, both algebras are deeply connected to each other and can be represented in the same space. The initial success of CFT was based on well-developed representation theory of the Virasoro algebra [9]. At the same time, ZF algebra approach seems to be more general since algebraic structures like (1.1)- (1.2) with matrices $\mathbf{R}$ and $\mathbf{W}$ depending on $\zeta_1\zeta_2^{-1}$, play crucial role in the two-dimensional Integrable models of both Statistical Mechanics ([10], [11], [12], [13], [14]) and Quantum Field Theory (see e.g. [15]) In the hierarchy of the solutions of YBE the constant solutions corresponding to algebras (1.1)- (1.2) are the simplest ones. They might be obtained from trigonometric and elliptic $\mathbf{R}$ and $\mathbf{W}$ matrices as a result of the well-known limiting procedure. Conversely, one can consider more complicated solutions of YBE as parametric deformations of the constant ones. It is reasonable to expect that ZF algebras corresponding to the trigonometric and elliptic matrices $\mathbf{R}$, $\mathbf{W}$ are deformations of the algebras of conformal vertex operators. In this way we run into the following questions:

1. *Is it possible to describe the representations of elliptic and trigonometric ZF algebras using the methods of CFT, or, more explicitly, can ZF algebras be realized in the direct sum of irreducible representations of some infinite-dimensional algebras generalizing Virasoro algebra?*
2. *What is the exact form of the commutation relations of these deformed Virasoro algebras and their geometrical and physical meaning?*

The present work is mainly devoted to studying of the first problem. We construct the representation of the elliptic deformation of the conformal vertex operator algebra (1.1). Our main idea is to deform in the appropriate manner the bosonization procedure developed in the works [2], [9], [16] for CFT. Let us recall that the central objects in the bosonization are screening operators. Indeed, to realize the irreducible representations of the Virasoro algebra in the Fock space, one needs to know only explicit bosonic realization of intertwining (screening) operators. It is remarkable that the explicit form of commutation relations of the Virasoro algebra is not really used. We describe a generalization of the conformal bosonization basing on deformed screening operators which was proposed in the work [17]. It allows one to get the explicit bosonic realization of the chiral vertex

operators which satisfy the commutation relations of the form (1.1) with elliptic $\mathbf{W}$ matrices. Quadratic associative algebras obtained in this approach seem to be deeply connected with algebras of vertex operators found in the works [18]. In particular, we expect that deformed Virasoro algebra is some reduction of the elliptic $sl(2)$ algebra as well as ordinary Virasoro algebra is a result of quantum Hamiltonian reduction of affine Kac-Moody algebra $\hat{sl}(2)$ [19].

The present work is prepared to be presented at the A.A.Belavin's seminar on CFT and Integrable Models at Chernogolovka. Its largest part is devoted to bosonization of CFT. In the simplest cases we re-examine why and when bosonization is still working. The material from the first and the third sections seems to be known to experts and it is contained (but sometimes in the hidden form) in the works [2], [9], [16]. Our aim here is just to emphasize those subtle moments which appeared to be crucial in the generalization of bosonization [17]. As soon as we will understand and correctly formulate conformal case, we will be able to construct the representation of the elliptic ZF algebra. In particular, we obtain integral representation for matrix elements of vertex operators generalizing Dotsenko-Fateev formulas for conformal blocks. For instance, in the simplest case of four-point function it is equivalent to the following integral representation of q-hypergeometrical function:

$$\begin{aligned} \int_C \frac{dz}{2\pi i} z^{c-1} \frac{(q^{\frac{1+a}{2}} z^{-1}; q)_\infty}{(q^{\frac{1-a}{2}} z^{-1}; q)_\infty} \frac{(q^{\frac{1+b}{2}} \zeta z; q)_\infty}{(q^{\frac{1-b}{2}} \zeta z; q)_\infty} = \\ = q^{\frac{c(1-a)}{2}} \frac{\Gamma_q(c+a)}{\Gamma_q(c+1) \Gamma_q(a)} F_q(a+c, b, c+1; q^{1-\frac{a+b}{2}} \zeta) . \end{aligned} \quad (1.4)$$

Notice, that matrix elements of deformed vertex operators are written in the form of ordinary contour integrals rather than Jackson's integrals in the bosonization scheme of the works [20]. We also consider trigonometric limit of elliptic construction which corresponds to Sin-Gordon model [17], [21], [22]. In this limit we show how to reconstruct the ZF algebra of vertex type from IRF ZF algebra [23].

Let us make some notational conventions.

- i) Throughout this work we will denote objects which have similar meaning by the same letter, distinguishing it by "prime" symbol '. It should not be confused with derivation symbol ∂ .
- ii) For the technical reason it is convenient for us to make ordering procedure in the exponential operators like $e^{i\phi}$ in the final step. There is no difference in this prescription with ordinary one in the conformal case, and it is not principal but useful in the deformed case. It will be explained in the section 4.

2. Preliminaries

2.1 Let $R_{ab}^{cd}(t)$ be numerical matrix depending on complex parameter t . The indexes a, b, c, d take value in the set of integer numbers. One can consider this matrix as an operator acting in tensor product $\mathcal{V} \otimes \mathcal{V}$, where $\mathcal{V}$ is finite dimensional linear space, which vectors are specified by indexes a . We will call by Yang-Baxter equation of vertex type the following algebraic equation [3] :

$$\begin{aligned} R_{a_1 a_2}^{c_1 c_2}(t_1 t_2^{-1}) R_{c_1 a_3}^{b_1 c_3}(t_1 t_3^{-1}) R_{c_2 c_3}^{b_2 b_3}(t_2 t_3^{-1}) = \\ = R_{a_1 c_3}^{c_1 b_3}(t_1 t_3^{-1}) R_{a_2 a_3}^{c_2 c_3}(t_2 t_3^{-1}) R_{c_1 c_2}^{b_1 b_2}(t_1 t_2^{-1}) . \end{aligned} \quad (2.1)$$

This equation play fundamental role in the 2 D Integrable Models of both Quantum Field Theory and Statistical Mechanics. At present a lot of its nontrivial solutions are found (see e.g. [24]). In this work we will be interested in the simplest one when the space $\mathcal{V}$ has the dimension equal 2. We will enumerate basic vectors of $\mathcal{V}$ by $\pm$. In 1972 Baxter found the following remarkable solution of (2.1) [25]:

$$\begin{aligned} R_{--}^{-}(t, p, q) &= \frac{\Theta_q(p^2 q^{\frac{1}{2}}) \Theta_q(t q^{\frac{1}{2}})}{\Theta_q(q^{\frac{1}{2}}) \Theta_q(p^2 t q^{\frac{1}{2}})} , \\ R_{+-}^{+}(t, p, q) &= R_{-+}^{-}(t, p, q) = p \frac{\Theta_q(p^2 q^{\frac{1}{2}}) \Theta_q(t)}{\Theta_q(q^{\frac{1}{2}}) \Theta_q(p^2 t)} , \\ R_{+-}^{-}(t, p, q) &= R_{-+}^{+}(t, p, q) = t^{\frac{1}{2}} \frac{\Theta_q(p^2) \Theta_q(t q^{\frac{1}{2}})}{\Theta_q(q^{\frac{1}{2}}) \Theta_q(p^2 t)} , \\ R_{++}^{-}(t, p, q) &= R_{++}^{-}(t, p, q) = p^{-1} t^{-\frac{1}{2}} q^{\frac{1}{4}} \frac{\Theta_q(p^2) \Theta_q(t)}{\Theta_q(q^{\frac{1}{2}}) \Theta_q(p^2 t q^{\frac{1}{2}})} , \end{aligned} \quad (2.2)$$

where

$$\Theta_q(s) = (q; q)_{\infty} (s; q)_{\infty} (q s^{-1}; q)_{\infty} ,$$

is the Jacobi elliptic function and we use the standard notation:

$$(z; q_1, \dots, q_k)_{\infty} = \prod_{n_1, \dots, n_k=0}^{\infty} (1 - z q_1^{n_1} \dots q_k^{n_k}) .$$

The R -matrix depends on two additional parameters p, q . In this work we suppose that parameter q is real and $0 \leq q < 1$. At the same time p will be complex number such that

$|p|^2 = 1$. In this case the matrix elements are real numbers if $|t|^2 = 1$. Together with the Yang-Baxter equation the matrix R_{ab}^{cd} satisfies also to so-called unitarity condition:

$$R_{a_1 a_2}^{b_1 b_2}(t) R_{b_1 b_2}^{c_1 c_2}(t^{-1}) = \delta_{a_1}^{c_1} \delta_{a_2}^{c_2}. \quad (2.3)$$

As it was pointed out by A.B. Zamolodchikov [6] the equations (2.1), (2.3) can be treated as correspondingly self consistency and associativity condition in the formal algebra:

$$Z_{a_1}(t_1) Z_{a_2}(t_2) = S_{a_1 a_2}^{b_1 b_2}(t_1 t_2^{-1}) Z_{b_2}(t_2) Z_{b_1}(t_1). \quad (2.4)$$

This algebra was also considered in the works of L.D. Faddeev [7] and we will call it as ZF algebra.

If the parameter $q \rightarrow 0$ the matrix extremely simplifies since $\Theta_q(t) \rightarrow 1 - t$ and nontrivial elements of the matrix $R_{ab}^{cd}(t, p)$ read:

$$\begin{aligned} R_{++}^{++}(t, p) &= R_{--}^{--}(t, p) = 1, \\ R_{+-}^{+-}(t, p) &= R_{-+}^{-+}(t, p) = p \frac{1-t}{1-p^2 t}, \\ R_{+-}^{+ -}(t, p) &= R_{-+}^{- +}(t, p) = t^{\frac{1}{2}} \frac{1-p^2}{1-p^2 t}. \end{aligned} \quad (2.5)$$

We can also get a solution of Yang-Baxter equation which does not depend on spectral parameter t by the following limiting procedure:

$$R_{ab}^{cd}(\sigma, p) = p^{\frac{\sigma}{2}} \lim_{L \rightarrow +\infty} e^{\frac{(\sigma-c)\sigma L}{4}} R_{ab}^{cd}(e^{\sigma L}, p), \quad (2.6)$$

here $\sigma = \pm$. Non zero elements of matrix $R_{ab}^{cd}(+, p)$ are defined by the relations

$$\begin{aligned} R_{++}^{++}(+, p) &= R_{--}^{--}(+, p) = p^{\frac{1}{2}}, \\ R_{+-}^{+-}(+, p) &= R_{-+}^{-+}(+, p) = p^{-\frac{1}{2}}, \\ R_{+-}^{+ -}(+, p) &= p^{-\frac{1}{2}}(p - p^{-1}). \end{aligned} \quad (2.7)$$

Note that matrices $R(+, p)$ and $R(-, p)$ are connected by the relation

$$R_{ab}^{cd}(-, p) = [R^{-1}]_{ba}^{dc}(+, p), \quad (2.8)$$

which can be considered as analogue of unitarity condition for constant solution of Yang-Baxter equation. The matrix (2.7) coincides with the fundamental R-matrix for quantum

algebra $U_p(sl(2))$ [26] . We will need some facts concerning this algebra, so let us recall them here.

2.2. The quantum universal enveloping algebra $U_p(sl(2))$ is an algebra on generators $X^\pm, T$ subject to the following relations:

$$\begin{aligned} X^+ X^- - p^2 X^- X^+ &= \frac{1 - T^2}{1 - p^{-2}} \\ TX^\pm &= p^{\mp 2} X^\pm T . \end{aligned} \quad (2.9)$$

It is well known that (2.9) has Hopf algebra structure. In particular, the comultiplication Δ is defined by

$$\Delta(X^\pm) = T \otimes X^\pm + X^\pm \otimes 1 , \Delta(T) = T \otimes T . \quad (2.10)$$

The parameter p is complex number. The most relevant cases for us will be those with $p = e^{i\pi \frac{\xi+1}{\xi}}$ or $e^{i\pi \frac{\xi}{\xi+1}}$ where ξ is a real irrational number greater then 1. In these cases the algebra $U_p(sl(2))$ admits the irreducible representations in the finite dimensional spaces $\mathcal{V}_l$ with the $\dim \mathcal{V}_l = l$. Construction of such representations is quite similar to the construction of irreducible representations of $sl(2)$ Lie algebra with the spin $j = \frac{l-1}{2}$ [27], [28] . The basic vectors $\mathbf{e}_l^m \in \mathcal{V}_l$ of representation are specified by number $m = -j, -j+1, \dots, j$ and conditions

$$\begin{aligned} T\mathbf{e}_l^m &= p^{-2m} \mathbf{e}_l^m , \\ X^- \mathbf{e}_l^m &= \mathbf{e}_l^{m-1} . \end{aligned} \quad (2.11)$$

The existence of Hopf structure means that tensor product $\mathcal{V}_{l_1} \otimes \mathcal{V}_{l_2}$ of two representations of $U_p(sl(2))$ would also carry the structure of representation of this algebra. Moreover, if ξ is an irrational number, then this representation turns to be completely reducible. Since the tensor product of two finite-dimensional irreducible representations can be represented as direct sum of irreducible representations, then one can write down the Clebsh-Gordon decomposition of the form:

$$\mathbf{e}_{2j_1+1}^{m_1} \otimes \mathbf{e}_{2j_2+1}^{m_2} = \sum_l \begin{pmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m_1 + m_2 \end{pmatrix}_p \mathbf{e}_{2j+1}^m . \quad (2.12)$$

In present work we will need the explicit form of the following Clebsh-Gordon coefficients:

$$\begin{aligned}
\left(\begin{array}{ccc} \frac{1}{2} & j & j + \frac{1}{2} \\ -\frac{1}{2} & m & m + \frac{1}{2} \end{array} \right)_p &= p^{-2j} \frac{[j + m + 1]_p}{[2j + 1]_p} , \\
\left(\begin{array}{ccc} \frac{1}{2} & j & j + \frac{1}{2} \\ -\frac{1}{2} & m & m - \frac{1}{2} \end{array} \right)_p &= \frac{p^{-j+m}}{[2j + 1]_p} , \\
\left(\begin{array}{ccc} \frac{1}{2} & j & j - \frac{1}{2} \\ -\frac{1}{2} & m & m + \frac{1}{2} \end{array} \right)_p &= (-1)^{2j} \frac{[j - m]_p}{[2j + 1]_p} , \\
\left(\begin{array}{ccc} \frac{1}{2} & j & j - \frac{1}{2} \\ -\frac{1}{2} & m & m - \frac{1}{2} \end{array} \right)_p &= (-1)^{2j+1} \frac{p^{-j+m-1}}{[2j + 1]_p} ,
\end{aligned} \tag{2.13}$$

here we use the notation $[x]_p = \frac{p^x - p^{-x}}{p - p^{-1}}$.

3. Free Fermions ZF algebra and $c = -2$ Virasoro algebra

In this section we will discuss ZF algebra generated by operators $Z_a(\zeta)$ ($a = \pm, \zeta \in \mathbf{C}$), satisfying the simple commutation relations

$$Z_a(\zeta_1)Z_b(\zeta_2) = -Z_b(\zeta_2)Z_a(\zeta_1), \quad \zeta_1 \neq \zeta_2 . \tag{3.1}$$

We consider its irreducible representations characterized by operator product expansions:

$$\begin{aligned}
Z_{\pm}(\zeta_2)Z_{\mp}(\zeta_1) &= \pm \frac{2\zeta_1\zeta_2}{(\zeta_2 - \zeta_1)^2} + O(1) , \\
Z_{\pm}(\zeta_2)Z_{\pm}(\zeta_1) &= O(1), \quad \zeta_1 \rightarrow \zeta_2 .
\end{aligned} \tag{3.2}$$

This algebra is well known in the physical literature as so called $b-c$ system ($Z_+ = b$, $Z_- = \partial c$). Its representations admit decompositions into direct sum of irreducible representations of the Virasoro algebra. Let us note now that the commutation relations (3.1) and operator product expansions (3.2) are invariant with respect to a linear transformation

$$Z_a[r] \rightarrow \mathbf{G}^{ab} Z_b[r] ,$$

if the determinant of $\mathbf{G}$ is equal to one. We will concentrate on the case when this symmetry is not broken and the representation space of the ZF algebra is classified by the $SL(2)$ symmetry together with Virasoro one. In this case Virasoro algebra has the central charge equal to -2. We will try to analyze these well-known results in the form which leaves room for generalization.

3.1. Irreducible representations of the free fermions algebra

The algebra (3.1) possesses two different types of representation denoted by π_Z^R (Ramond) and π_Z^{NS} (Neveu-Schwartz) according to type of boundary conditions imposed on the generators $Z_a(\zeta)$:

$$\begin{aligned} R : \quad Z_a(e^{2\pi i}\zeta) &= Z_a(\zeta) , \\ NS : \quad Z_a(e^{2\pi i}\zeta) &= -Z_a(\zeta) . \end{aligned} \quad (3.3)$$

For this reason, Z_a have the following decomposition in Laurent series:

$$Z_a(\zeta) = \sum_{r \neq 0} Z_a[r] \zeta^{-r} , \quad (3.4)$$

where $r \in \mathbf{Z}$ for R sector and $r \in \mathbf{Z} + \frac{1}{2}$ for NS one. It follows from (3.1)- (3.2) , that modes $Z_a[r]$ obey the anticommutation relations:

$$\{Z_a[r], Z_b[m]\}_+ = 2 \, r \, a \, \delta_{a+b,0} \delta_{r+m,0} . \quad (3.5)$$

The spaces π_Z^R and π_Z^{NS} are defined as Fock modules of fermionic algebra (3.5) created by the operators $Z_a[r]$, $r < 0$ under the action on the corresponding R- and NS-vacuum states. The vacuum states are specified by the condition that they are annihilated by any operator $Z_a[r]$ with $r > 0$. We assume that R- and NS- vacuum states are scalars with respect to the $SL(2)$ transformations. Then $SL(2)$ structure of the spaces $\pi_Z^{R,NS}$ is uniquely determined by the condition that ZF operators are arranged in the fundamental $SL(2)$ doublet.

The next fact which we will need further is that the dual space $\pi_Z^{R,NS\star}$ also admits the structure of representation of the algebra (3.5). It goes as follows. Given a linear space π_Z its dual $\pi_Z^\star$ is the set of linear maps $\pi_Z^\star : \pi_Z \rightarrow \mathbf{C}$ determined by action $\mathbf{u}^\star \longrightarrow \langle \mathbf{u}^\star, \mathbf{u} \rangle$. Choose the dual basis by the canonical pairing $\langle \mathbf{u}_j^\star, \mathbf{u}_k \rangle = \delta_{jk}$. Since algebra (3.5) admits the anti-involution

$$\mathcal{A} \{Z_a[r]\} = Z_{-a}[-r] , \quad (3.6)$$

then the spaces $\pi_Z^\star = \pi_Z^{R,NS\star}$ can be endowed by the structure of representation of ZF algebra through the formula

$$\langle \mathbf{u}_1^\star Z_a^\star[n], \mathbf{u}_2 \rangle = \langle \mathbf{u}_1^\star, \mathcal{A} \{Z_a[n]\} \mathbf{u}_2 \rangle . \quad (3.7)$$

3.2. Bosonization of fermionic ZF algebra

Now we want to show how the representations of the ZF algebra (3.1)-(3.2) can be realized in direct sums of boson Fock modules. We start with the space π_Z^R . Let $\{b_n, P, Q \mid n \in \mathbf{Z}/\{0\}\}$ be a set of operators satisfying the commutation relations:

$$\begin{aligned} [b_m, b_n] &= m \delta_{m+n,0} , \\ [P, Q] &= -i, \quad [P, b_n] = [Q, b_n] = 0 . \end{aligned} \quad (3.8)$$

The bosonic Fock module $\mathcal{F}_p$ is generated by acting of creation operators $b_{-n}, n > 0$ on the highest vector $\mathbf{f}_p$: $b_n \mathbf{f}_p = 0, \quad n > 0$; $P \mathbf{f}_p = p \mathbf{f}_p$. In the direct sum of the Fock modules $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}$ action of the following operators is well-defined:

$$\begin{aligned} X &= \oint \frac{dz}{2\pi i z} e^{-2i\phi(z)}, \\ X' &= \oint \frac{dz}{2\pi i z} e^{i\phi(z)}, \end{aligned} \quad (3.9)$$

where the integration contours are around the zero and

$$\phi(z) = Q - i P \ln z + \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} \frac{b_m}{i m} z^{-m} .$$

It is easy to see that operator X' is nilpotent one, $X'^2 = 0$. Let us define the following spaces:

$$\begin{aligned} \mathcal{F}_{k-\frac{1}{2}}^+ &= \text{Ker}_{\mathcal{F}_{k-\frac{1}{2}}} [X'] , \\ \mathcal{F}_{k-\frac{1}{2}}^- &= \mathcal{F}_{k-\frac{1}{2}} / \mathcal{F}_{k-\frac{1}{2}}^+ . \end{aligned} \quad (3.10)$$

Since the operators X, X', P obey the commutation relations:

$$[X, X'] = 0, \quad [P, X] = -2X, \quad [P, X'] = X' , \quad (3.11)$$

then they act as

$$\begin{aligned} \mathcal{F}_{k+\frac{1}{2}}^\pm &\xrightarrow{X} \mathcal{F}_{k-\frac{3}{2}}^\pm , \\ \mathcal{F}_{k-\frac{1}{2}}^- &\xrightarrow{X'} \mathcal{F}_{k+\frac{1}{2}}^+ \xrightarrow{X'} 0 . \end{aligned} \quad (3.12)$$

In the simple case we are considering now, it is not hard to guess the expressions of $Z_a(\zeta)$ through the field ϕ . Finally, the bosonization of π_Z^R is described by the

Proposition 3.1 *The map $\pi_Z^R \rightarrow \bigoplus_{k \in \mathbb{Z}} \mathcal{F}_{k-\frac{1}{2}}^+$ given by the identification of the R -vacuum state of π_Z^R with vector $\mathbf{f}_{\frac{1}{2}}$ and bosonization rules*

$$\begin{aligned} Z_+(\zeta) &= e^{i\phi(\zeta)} , \\ Z_-(\zeta) &= [X, Z_+(\zeta)] = 2\zeta \partial_\zeta e^{-i\phi(\zeta)} \end{aligned} \quad (3.13)$$

is an isomorphism of modules.

Note that it follows from the formulae (3.11) that operators X and $H = P - \frac{1}{2}$ can be considered as generators of the Borel subalgebra of $sl(2)$ algebra acting in the space π_Z^R .

Basing on the proposition 3.1, one can treat π_Z^R as a direct sum of Virasoro algebra modules. The crucial observation here is that the subspaces $\text{Ker}_{\mathcal{F}_{l-\frac{1}{2}}^+} [X^l], l > 0$ can be endowed by the structure of irreducible representation of the Virasoro algebra. To explain this important property let us introduce $SL(2)$ scalar field $T(\zeta)$ by the formula:

$$-4 \zeta^2 T(\zeta) = Z_+(\zeta) Z_-(\zeta) - Z_-(\zeta) Z_+(\zeta) . \quad (3.14)$$

In terms of bosonic field ϕ it reads

$$2 T(\zeta) = f^2 - \partial_\zeta f , \quad (3.15)$$

where $f = i\partial_\zeta \phi(\zeta) - \frac{1}{2\zeta}$. It is easy to check that $L_n : T(\zeta) = \sum_{n \in \mathbb{Z}} L_n \zeta^{-n-2}$ generate the Virasoro algebra Vir_c

$$[L_n, L_m] = (n - m) L_{n+m} + \frac{c}{12} (n^3 - n) \quad (3.16)$$

with $c = -2$. Let us denote by $\mathcal{L}_{l-\frac{1}{2}}, l > 0$ the irreducible Verma module with highest weight

$$\Delta_{l-\frac{1}{2}} = \frac{l(l-1)}{2} \quad (3.17)$$

built upon highest vector $\mathbf{v}_{l-\frac{1}{2}}$. If generators of the Virasoro algebra are realized as (3.15) then any highest weight vector $\mathbf{f}_{l-\frac{1}{2}}$ of Fock space turns to be the highest weight vector of Verma module of Virasoro algebra with highest weight (3.17). Moreover, consider in bosonic space π_Z^R non-zero vectors which can be obtained from $\mathbf{f}_{l-\frac{1}{2}}$ by the action of operator $X^k, k = 0, 1, \dots$. We claim that generators L_n with $n > 0$ annihilate any such vector while L_0 acts by multiplication on constant $\Delta_{l-\frac{1}{2}}$. It follows from the fact that generators (3.15) commute with operator X as it can be easily checked.

Let us consider formal tensor product of irreducible Virasoro algebra module $\mathcal{L}_{l-\frac{1}{2}}$ and l -dimensional $sl(2)$ irreducible representation $\mathcal{V}_l^+$ with basic vectors $\mathbf{e}_l^{m+}$, where $m = -j, \dots, j$ and $l = 2j + 1$, $j = 0, \frac{1}{2}, 1, \dots$. We will denote it as $\mathcal{L}_{l-\frac{1}{2}} \otimes \mathcal{V}_l^+$. The meaning of the additional index "+" will be clarified later. Standard arguments [9] possess to prove the following proposition describing Virasoro structure of the space π_Z^R (fig. 1):

Proposition 3.2

The map $\pi_Z^R \rightarrow \oplus_{l=1}^{\infty} \mathcal{L}_{l-\frac{1}{2}} \otimes \mathcal{V}_l^+$ given by correspondence $X^{j-m} \mathbf{f}_{l-\frac{1}{2}} \rightarrow \mathbf{v}_{l-\frac{1}{2}} \otimes \mathbf{e}_l^{m+}$ ($l = 2j + 1$) and bosonization rule (3.15) is an isomorphism of modules of the Virasoro algebra with

i)

$$\text{Ker}_{\mathcal{F}_{l-\frac{1}{2}}^+} [X^l] \cong \mathcal{L}_{l-\frac{1}{2}}, l > 0; \quad (3.18)$$

ii)

$$\mathcal{F}_{2m+\frac{1}{2}}^+ \cong \oplus_{k=0}^{\infty} \mathcal{L}_{2|m|+2k+\frac{1}{2}} \otimes \mathbf{e}_{2|m|+2k+1}^{m+}, m \in \frac{1}{2}\mathbf{Z}. \quad (3.19)$$




Fig. 1: The Virasoro structure of the space $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^+$

Notice, that such as operators L_n and X' commute, then the space $\mathcal{F}_{k-\frac{1}{2}}^-$ has the similar Virasoro structure as $\mathcal{F}_{k+\frac{1}{2}}^+$:

$$\mathcal{F}_{2m-\frac{1}{2}}^- \cong \oplus_{k=0}^{\infty} \mathcal{L}_{2|m|+2k+\frac{1}{2}} \otimes \mathbf{e}_{2|m|+2k+1}^{m-}, m \in \frac{1}{2}\mathbf{Z}, \quad (3.20)$$

where we denoted by $\mathbf{e}_l^{m-}$ the basis vectors in l -dimensional space $\mathcal{V}_l^-$. Note that vector $\mathbf{v}_{l-\frac{1}{2}} \otimes \mathbf{e}_l^{-m,-}$ where $\mathbf{v}_{l-\frac{1}{2}}$ is highest vectors in Virasoro module is identified with the state from $\mathcal{F}_{-2m-\frac{1}{2}}^-$ which is the proimage of $X^{j+m} \mathbf{f}_{-l+\frac{1}{2}}$ with respect to the action of operator X' . In particular $\mathbf{v}_{l-\frac{1}{2}} \otimes \mathbf{e}_l^{-j,-} \equiv \mathbf{f}_{-l+\frac{1}{2}} \mod Ker X'$. The difference which appears in this case is in the fact that irreducible Virasoro algebra modules are identified with factor spaces rather than subspaces of Fock space.

3.3. Scalar product in Fock space.

Now we turn to the bosonization of dual representation $\pi_Z^{R\star}$ of ZF algebra. This problem is closely related with the proper choice of the scalar product in the direct sum of Fock spaces $\oplus_{k \in \mathbb{Z}} \mathcal{F}_{k-\frac{1}{2}}$. The obvious guess is

$$(\mathbf{u}_1 b_n, \mathbf{u}_2)_0 = -(\mathbf{u}_1, b_{-n} \mathbf{u}_2)_0, (\mathbf{u}_1 P, \mathbf{u}_2)_0 = -(\mathbf{u}_1, P \mathbf{u}_2)_0. \quad (3.21)$$

However, it does not conform with conjugation condition (3.7) for operators Z_a . In thinking about this problem, it is rather natural to introduce independent bosonic representation for $\pi_Z^{R\star}$ and then try to identify it with some subspaces in the Fock space $\oplus_{k \in \mathbb{Z}} \mathcal{F}_{k-\frac{1}{2}}$. To do this, we define new set of generators $\{b_n^*, P^*, Q^* | n = \pm 1, \pm 2, \dots\}$ with the commutation relations as in (3.8). Let analogues for integral operators (3.9) be

$$\begin{aligned} X'^{\star} &= \oint \frac{dz}{2\pi i z} e^{i\phi^*(z)}, \\ X^{\star} &= \oint \frac{dz}{2\pi i z} e^{-2i\phi^*(z)}. \end{aligned} \quad (3.22)$$

Repeating step by step the analysis above, one can describe the bosonization of the representation $\pi_Z^{R\star}$ and introduce Virasoro algebra structure in it. The formulae for this case are rather evident. In particular $\pi_Z^{R\star} \cong \oplus_{k \in \mathbb{Z}} \mathcal{F}_{k-\frac{1}{2}}^{\star+}$, where we denote by $\mathcal{F}_{k-\frac{1}{2}}^{\star+}$ the kernel of the operator $X'^{\star} : \mathcal{F}_{k-\frac{1}{2}}^{\star} \rightarrow \mathcal{F}_{k+\frac{1}{2}}^{\star}$. And any space $\mathcal{F}_{k+\frac{1}{2}}^{\star+}$ turns to be isomorphic to direct sum of irreducible Virasoro algebra modules :

$$\mathcal{F}_{2m+\frac{1}{2}}^{\star+} \cong \oplus_{k=0}^{\infty} \mathcal{L}_{2|m|+2k+\frac{1}{2}}^{\star} \otimes \mathbf{e}_{2|m|+2k+1}^{\star m}. \quad (3.23)$$

Here the states $X^{\star j-m} \mathbf{f}_{l-\frac{1}{2}}^{\star}$ ($l = 2j+1$) correspond to states $\mathbf{v}_{l-\frac{1}{2}}^{\star} \otimes \mathbf{e}_l^{\star m}$. Notice, that the symbol $\star$ in the $\mathcal{L}_{l-\frac{1}{2}}^{\star}$ is used in order to emphasize the fact that generators of Virasoro algebra are built from fields $\phi^{\star}$ rather than from ϕ 's. To proceed further, let us note, that the decompositions (3.20) , (3.23) of spaces $\pi_Z^{R\star}$ and $\oplus_{k \in \mathbb{Z}} \mathcal{F}_{k-\frac{1}{2}}^-$ include the identical

sets of irreducible representations of the Virasoro algebra. Indeed, any two representations $\mathcal{L}_{l-\frac{1}{2}}$ and $\mathcal{L}_{l-\frac{1}{2}}^*$ are isomorphic as $c = -2$ Virasoro algebra representations since they have the same highest weight $\Delta_{l-\frac{1}{2}}$. In the bosonic realization this means that one can identify the Virasoro algebra generators.¹ To extend isomorphism of Virasoro algebra modules on the isomorphism of $\pi_Z^{R\star}$ and $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^-$ one needs establish the correspondence between vectors in $\mathcal{V}_l^*$ and $\mathcal{V}_l^-$. Let us identify vectors $\mathbf{e}_l^{m\star} \in \mathcal{V}_l^*$ and $c_m^j \mathbf{e}_l^{-m,-} \in \mathcal{V}_l^-$. The constants c_m^j can be fixed by the condition that the operators X, X^* and $H = -P - \frac{1}{2}$ acting in the space $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^- \equiv \oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^{*\star+}$ generate the $sl(2)$ Lie algebra.

$$[X, X^*] = H, \quad [H, X] = 2X, \quad [H, X^*] = -2X^* . \quad (3.24)$$

Then we will have $c_m^j = \frac{(j-m)!(2j)!}{(j+m)!}$. Hence, we conclude that the representation $\pi_Z^{R\star}$ can be realized as following:

$$\pi_Z^{R\star} = \oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^- . \quad (3.25)$$

Note that due to this identification, an action of the operator X^* in the space $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}^-$ can be uniquely specified by the condition that the following diagram is commutative:

$$\begin{array}{ccccc} & & X' & & \\ & \mathcal{F}_{k-\frac{3}{2}}^- & \longrightarrow & \mathcal{F}_{k-\frac{1}{2}}^+ & \\ X^* & \downarrow & X' & \downarrow & X^* \\ & \mathcal{F}_{k+\frac{1}{2}}^- & \longrightarrow & \mathcal{F}_{k+\frac{3}{2}}^+ & \end{array} . \quad (3.26)$$

Now we are able to describe total symmetry algebra $Symm$ which acts in the space $\pi_Z^R \oplus \pi_Z^{R\star}$. First, it includes infinite dimensional Virasoro algebra with central charge $c = -2$. Second, it contains $sl(2)$ subalgebra. Finally, the operators X' and X'^* generate the Clifford subalgebra:

$$\{X', X'^*\}_+ = 1, \quad X'^2 = [X'^*]^2 = 0 . \quad (3.27)$$

Note that the operators X', X'^* commute with Virasoro generators and X, X^* but do not commute with H . The decomposition of the space $\pi_Z^R \oplus \pi_Z^{R\star}$ into direct sum of irreducible representations of the algebra $Symm$ has the form:

$$\pi_Z^R \oplus \pi_Z^{R\star} = \oplus_{l=0}^{\infty} \mathcal{L}_{l-\frac{1}{2}} \otimes \mathcal{V}_l \otimes \mathcal{V}_2' , \quad (3.28)$$

¹ Because of this identification one can consider fields $i\partial_\zeta \phi - \frac{1}{2\zeta}$ and $i\partial_\zeta \phi^* - \frac{1}{2\zeta}$ as different solutions of quantum version of the Riccati equation (3.15) [4] .

where we denoted the l -dimensional irreducible representation of algebra $su(2)$ as $\mathcal{V}_l$ and $\mathcal{V}'_2$ is two dimensional representation of Clifford algebra (3.27). Decomposition (3.28) makes clear the symbolic notations introduced earlier. Namely, vectors $\mathbf{e}_l^{m\pm}$ are basic vectors in the irreducible l -dimensional representations of $sl(2)$ algebra with spin $j = \frac{l-1}{2}$ and momentum projection m . Any pair of vectors $\mathbf{e}_l^{m-}$ and $\mathbf{e}_l^{m+}$ is arranged in doublet of the Clifford algebra.

Let us discuss now the Hilbert structure of the space $\pi_Z^R \oplus \pi_Z^{R*}$. This space has a canonical scalar product $(\mathbf{u}_1, \mathbf{u}_2)$ induced by condition: $(\mathbf{u}_1, \mathbf{u}_2) = (\mathbf{u}_1^*, \mathbf{u}_2^*) = 0$, $(\mathbf{u}_1^*, \mathbf{u}_2) = (\mathbf{u}_1, \mathbf{u}_2^*)^* = \langle \mathbf{u}_1^*, \mathbf{u}_2 \rangle$ for any $\mathbf{u}_{1,2} \in \pi_Z^R$, $\mathbf{u}_{1,2}^* \in \pi_Z^{R*}$. From the other side such scalar product is equivalently described as following:

$$(\mathbf{e}_{2j'+1}^n{}^b, \mathbf{e}_{2j+1}^m{}^a) = (-1)^{j+m} \frac{2^{2j}(2j)!}{2j+1} \delta_{j,j'} \delta_{m+n,0} \delta_{a+b,0} \quad (3.29)$$

and the scalar product in the irreducible representations of Virasoro algebra is defined by the conjugation conditions

$$(\mathbf{v}_1 L_n, \mathbf{v}_2) = (\mathbf{v}_1, L_{-n} \mathbf{v}_2) . \quad (3.30)$$

We will also assume that the scalar product of the highest Virasoro vectors have the form

$$(\mathbf{v}_{l_1-\frac{1}{2}}^*, \mathbf{v}_{l_2-\frac{1}{2}}) = l_1 \delta_{l_1, l_2} . \quad (3.31)$$

Notice that the scalar product (3.29)- (3.30) is drastically different from (3.21).

3.4. NS sector.

Now let us shortly describe the bosonization of NS sector of representation of ZF algebra. The consideration here is quite similar as it have been done for R-sector and we present only results. The space π_Z^{NS} can be realized as a following direct sum of Fock modules:

$$\pi_Z^{NS} = \oplus_{k \in \mathbb{Z}} \mathcal{F}_k , \quad (3.32)$$

where NS-vacuum is identified with $\mathbf{f}_0$. The expression for operators $Z_a(\zeta)$ in terms of bosonic field ϕ is given by formula (3.13) again. The main difference here in comparison with Ramond sector case is that operator X' does not act in NS sector. It happens since the integration contour in the definition of X' is not closed.

The states in the Fock module $\mathcal{F}_k$ can be classified with respect to the action of direct product of Virasoro and $sl(2)$ algebras:

$$\mathcal{F}_{2m} \cong \oplus_{k=0}^{\infty} \mathcal{L}_{2|m|+2k} \otimes \mathbf{e}_{2|m|+2k+1}^m, \quad m \in \frac{1}{2}\mathbf{Z}, \quad (3.33)$$

where we denoted by $\mathcal{L}_l$, $l > 0$ irreducible modules of Virasoro algebra built upon highest vector $\mathbf{v}_l$ with highest weight

$$\Delta_{l-1} = \frac{(l-1)^2}{2} - \frac{1}{8}. \quad (3.34)$$

Vectors $\mathbf{e}_l^m$ are basic vectors in the l -dimensional irreducible representation of the $sl(2)$ -algebra. Again, the correspondence between representations is given by the maps of vectors $X^{j-m} \mathbf{f}_l \rightarrow \mathbf{v}_l \otimes \mathbf{e}_l^m$ and bosonization rules for generators of Virasoro algebra and $sl(2)$ algebra. The scalar product in the space π_Z^{NS} can be introduced analogously as for R-sector. Notice, that in this case the space π_Z^{NS} can be considered as selfdual representation of ZF algebra:

$$\pi_Z^{NS} = \pi_Z^{NS*}.$$

3.5. Vertex operators for Vir_{-2} algebra.

Now, we wish to find an exact meaning of the operators $Z_{\pm}$ in terms of $Symm$ algebra. Before going on, let us explain the situation on which we will focus. In case under consideration the algebra $Symm$ turns to be a direct product of infinite-dimensional Virasoro algebra and finite algebra. We expect, this statement is general in the sense that it is possible to associate for any ZF algebra another algebra which would be a direct product of two parts, infinite-dimensional (like Virasoro) and finite ones. We will demonstrate by executing simple example (3.1) - (3.2) that these operators can be identified with vertex operators for the total symmetry algebra $Symm$. Keeping in the mind the decompositions (3.28) it is rather natural to study the action of the operators $Z_{\pm}$ on the Virasoro states separately from the action on the finite-dimensional part of the representation space. The operators $Z_{\pm}$ have the spins $\frac{1}{2}$ and momentum projections $\pm\frac{1}{2}$ with respect to $sl(2)$ algebra. Then, if $\mathbf{e}_l^m$ are basic vectors in irreducible representations $\mathcal{V}_l$ and $\mathbf{v} \in \mathcal{L}_p$, where $p = l - \frac{1}{2}, p = l - 1$ correspondingly for R and NS sectors, the operators $Z_a(\zeta)$ in the representations π_Z^R and π_Z^{NS} can be represented in the form:

$$Z_a(\zeta) \mathbf{v} \otimes \mathbf{e}_l^m = \sum_{b=\pm 1} \begin{pmatrix} \frac{1}{2} & j & j + \frac{b}{2} \\ \frac{a}{2} & m & m + \frac{a}{2} \end{pmatrix} \Phi_{21}^{p+b, p}(\zeta) \mathbf{v} \otimes \mathbf{e}_{l+b}^{m+\frac{a}{2}}, \quad (3.35)$$

where $l = 2j + 1$. The numerical coefficients in this formula coincide with the Clebsch-Gordan coefficients for $sl(2)$ algebra. At the same time the operators Φ_{21} act as following:

$$\Phi_{21}^{p\pm 1}(\zeta) : \mathcal{L}_{\Delta_p} \xrightarrow{\Phi_{21}} \mathcal{L}_{\Delta_{p\pm 1}} \otimes \mathbf{C}[\zeta] \zeta^{\Delta_{p\pm 1} - \Delta_p} , \quad (3.36)$$

where $\mathbf{C}[\zeta]$ denotes the Laurent series in ζ . Using operators $Z^\star \in \text{End}(\pi_Z^{R, NS^\star})$ one can define the operators $\Phi_{21}^\star$ by similar expressions. It follows from the formula (3.7) that the operators Φ_{21} and $\Phi_{21}^\star$ (3.36) satisfy the relation:

$$(\mathbf{v}_1^\star \Phi_{21}^{p\pm 1}(\zeta), \mathbf{v}_2) = (\mathbf{v}_1^\star, \Phi_{21}^{p\pm 1}(\zeta) \mathbf{v}_2) , \quad (3.37)$$

where $\mathbf{v}_1^\star \in \mathcal{L}_p^\star$ and $\mathbf{v}_2 \in \mathcal{L}_{p\pm 1}$.

We claim that the operators (3.36) are vertex operators for the Virasoro algebra. Indeed, using the bosonization (3.13) and conjugation condition (3.37) one can show that they satisfy the following commutation relations with Virasoro algebra generators L_n :

$$[L_n, \Phi_{21}^\pm(\zeta)] = \zeta^{n+1} \partial_\zeta \Phi_{21}^\pm(\zeta) + \Delta_{21} n \zeta^n \Phi_{21}^\pm(\zeta) , \quad (3.38)$$

where $\Delta_{21} = 1$. Here and after we will use the short notation $\Phi_{21}^\pm$ for operators $\Phi_{21}^{p\pm 1}$.

Thus, ZF algebra appeared to be naturally connected with chiral primary operators of Virasoro algebra. The last objects are of a great important for our next constructions. It is well-known [1], [2] that chiral primary operators are uniquely determined by the commutation relations with L_n , intertwining property and conjugation condition. From now on in this section let us assume that symbol $\mathcal{L}_p$ denotes irreducible representation of $c = -2$ Virasoro algebra with highest weight

$$\Delta_p = \frac{p^2}{2} - \frac{1}{8} ,$$

where $p \in \mathbf{R}$ rather than just integer or half-integer. This generalization is very useful, since, considering general case we will be able separate general features of the bosonic construction from the specific properties of $SL(2)$ -invariant fermion model (3.1)-(3.2).

Usually, the special attention is paid to the studying of the properties of two chiral primary operators [1], [2]. The first of these operators is given by Φ_{21} with properties (3.36) -(3.38) while the second can be introduced by the following conditions:

1. Intertwining properties

$$\Phi_{12}^{p \mp \frac{1}{2} p}(\zeta) : \mathcal{L}_p \xrightarrow{\Phi_{12}} \mathcal{L}_{p \mp \frac{1}{2}} \otimes \mathbf{C}[\zeta] \zeta^{\Delta_{p \mp \frac{1}{2}} - \Delta_p} . \quad (3.39)$$

2. Conjugation conditions:

$$(\mathbf{v}_1^* \Phi_{12}^{p \mp \frac{1}{2} p}(\zeta), \mathbf{v}_2) = (\mathbf{v}_1^*, \Phi_{12}^{p \mp \frac{1}{2} p}(\zeta), \mathbf{v}_2) . \quad (3.40)$$

3. Commutation relations with Virasoro algebra generators:

$$[L_n, \Phi_{12}^{\pm}(\zeta)] = \zeta^{n+1} \partial_{\zeta} \Phi_{12}^{\pm}(\zeta) + \Delta_{12} n \zeta^n \Phi_{12}^{\pm}(\zeta) , \quad (3.41)$$

with $\Delta_{12} = -\frac{1}{8}$.

Again, short notations $\Phi_{12}^{\pm}$ stand for operators $\Phi_{12}^{p \mp \frac{1}{2} p}$ correspondingly. The properties (3.36)-(3.38) and (3.39)-(3.41) describe chiral primary operators Φ_{21} and Φ_{12} uniquely up to multiple constant. This means that basing only on these formulae one can reconstruct matrix elements of any combination of such operators. Indeed, as it was shown by Belavin, Polyakov and Zamolodchikov [1], due to formulae (3.38) - (3.41) matrix elements of operators built from Φ_{21} and Φ_{12} turn to be solutions of certain linear differential equations. For instance, the functions

$$\begin{aligned} G_p^{\pm}(\zeta_1 \zeta_2^{-1}) &= (\mathbf{v}_p^*, \Phi_{21}^{\pm}(\zeta_2) \Phi_{21}^{\mp}(\zeta_1) \mathbf{v}_p) , \\ G_p'^{\pm}(\zeta_1 \zeta_2^{-1}) &= (\mathbf{v}_p^*, \Phi_{12}^{\pm}(\zeta_2) \Phi_{12}^{\mp}(\zeta_1) \mathbf{v}_p) , \end{aligned} \quad (3.42)$$

where $\mathbf{v}_p$ is highest vector in the irreducible Virasoro module $\mathcal{L}_p$, satisfy to the second order ordinary differential equations:

$$L_{\Delta_{21}} G_p^{\pm}(\zeta) = 0 , \quad L_{\Delta_{12}} G_p'^{\pm}(\zeta) = 0 . \quad (3.43)$$

The explicit form of linear differential operator L_{Δ} is given by:

$$L_{\Delta} = \left\{ \frac{3}{2(2\Delta + 1)} \partial_{\zeta}^2 + \left(\frac{1}{\zeta} + \frac{1}{\zeta - 1} \right) \partial_{\zeta} - \frac{\Delta_p}{\zeta^2} - \frac{\Delta}{(1 - \zeta)^2} + \frac{2\Delta}{\zeta(\zeta - 1)} \right\} \zeta^{-\Delta} . \quad (3.44)$$

According to this formula the functions (3.42) have the following asymptotics under $\zeta \rightarrow 0$:

$$G_p^{\pm}(\zeta) = O(\zeta^{\frac{1}{2} \mp p}) , \quad G_p'^{\pm}(\zeta) = O(\zeta^{\frac{1}{8} \pm \frac{p}{2}}) .$$

Up to normalization, the solutions of the equations (3.43) with boundary conditions above can be easily expressed through hypergeometric functions $F(a, b, c; \zeta)$ as:

$$\begin{aligned} G_p^\pm(\zeta) &= \zeta^{\frac{1}{2} \mp p} \frac{1 \mp 2p + \zeta(1 \pm 2p)}{(1 - \zeta)^2} , \\ G_p'^\pm(\zeta) &= \frac{\Gamma(\pm p + \frac{1}{2})}{\pi^{\frac{1}{2}} \Gamma(\pm p + 1)} \zeta^{\frac{1}{8} \pm \frac{p}{2}} (1 - \zeta)^{\frac{1}{4}} F(\pm p + \frac{1}{2}, \frac{1}{2}, \pm p + 1; \zeta) . \end{aligned} \quad (3.45)$$

Let us now take numbers $p \neq 0, \frac{1}{2}, 1, \dots$. Knowing explicit formulae (3.45) one can find the commutation relations of the algebra of chiral primary operators. It can be done by using the relation:

$$\begin{aligned} F(a, b, c; z) &= \frac{\Gamma(c) \Gamma(b - a)}{\Gamma(b) \Gamma(c - a)} (-z)^{-a} F(a, a - c + 1, a - b + 1; z^{-1}) \\ &+ \frac{\Gamma(c) \Gamma(a - b)}{\Gamma(a) \Gamma(c - b)} (-z)^{-b} F(b, b - c + 1, b - a + 1; z^{-1}) . \end{aligned} \quad (3.46)$$

The straightforward calculation ensures that operators $\Phi_{1,2}$ generate ZF algebra of IRF type:

$$\Phi_{12}^a(\zeta_1) \Phi_{12}^b(\zeta_2)|_{\mathcal{L}_p} = \sum_{c+d=a+b} \mathbf{W} \left[\begin{matrix} p + \frac{a+b}{2} & p + \frac{c}{2} \\ p + \frac{b}{2} & p \end{matrix} \middle| \sigma_{12} \right] \Phi_{12}^d(\zeta_2) \Phi_{12}^c(\zeta_1)|_{\mathcal{L}_p} . \quad (3.47)$$

The precise form of nontrivial elements of matrix $\mathbf{W}$ is the following:

$$\begin{aligned} \mathbf{W} \left[\begin{matrix} p \pm 1 & p \pm \frac{1}{2} \\ p \pm \frac{1}{2} & p \end{matrix} \middle| \sigma \right] &= e^{\frac{i\pi\sigma}{4}} , \\ \mathbf{W} \left[\begin{matrix} p & p \pm \frac{1}{2} \\ p \pm \frac{1}{2} & p \end{matrix} \middle| \sigma \right] &= \mp \frac{e^{-\frac{i\pi\sigma(1 \pm 4p)}{4}}}{\sin \pi p} , \\ \mathbf{W} \left[\begin{matrix} p & p \mp \frac{1}{2} \\ p \pm \frac{1}{2} & p \end{matrix} \middle| z \right] &= \pm e^{-\frac{\pi i \sigma}{4}} \cot \pi p . \end{aligned} \quad (3.48)$$

The commutation relations (3.47) should be considered as a rule of analytical continuation of functions from region $|\zeta_1| > |\zeta_2|$ to $|\zeta_1| < |\zeta_2|$ along the paths $C_{\sigma_{12}}, \sigma_{12} = \pm$.

Therefore, the dependence of the matrix $\mathbf{W}$ on the variable z appears only in the choice of the contour of the analytical continuation. The commutation relations of operators Φ_{21} are more simple. It hold for any value of p :

$$\Phi_{21}^a(\zeta_1) \Phi_{21}^b(\zeta_2) = -\Phi_{21}^b(\zeta_2) \Phi_{21}^a(\zeta_1) . \quad (3.49)$$




Fig. 2: Two contours of analytical continuations $C_{\pm}$ of functions $G_p^{\prime\pm}(\zeta)$.

If number p is integer or half-integer, then commutation relations of chiral primaries Φ_{12}^a become undetermined and algebra of chiral primaries would be meaningless. The technical reason for this inconsistency is in the fact that under half-integer values of p the hypergeometric function in $G_p^{\prime\pm}(\zeta)$ degenerates. Such phenomena is well-known. It appears if one study degenerated modules of Virasoro algebra with rational central charge.² To avoid additional complications we omit such cases, considering either special value of central charge c and modules in generic position (further in this section and in section 4), or degenerated modules but for generic central charge (as in section 5 below).

In spite of failure with algebra of chiral primaries, the model with $c = -2$ and $p = 0, \frac{1}{2}, 1, \dots$ elaborated above has several attractive features. First, the algebra of operators Φ_{21} is still the same as for the generic p . The fermionic representation of these operators drastically simplify the analysis of the construction. Second, in this model we have explicit bosonic realization of finite-dimensional subalgebra in the symmetry algebra which is given by operators X, X', P . And finally, such model gives us example how to find the bosonization of chiral primary operators. One can expect that form of bosonic realization of chiral operators should be the same, no matter whether the action of intertwining operators X, X' is defined or not.

3.6. Bosonization of vertex operators

Definitions of chiral primaries do not assume bosonization procedure. However, we will find bosonic realization of such operators because it admits direct generalization for

² In certain cases (minimal models) it is possible to reduce the space of states such that chiral primaries would be correctly defined and generate associative algebra [29].

less trivial cases. Since chiral primary operators do not depend on the realization of $\mathcal{L}_p$, then its matrix elements are the same for any irreducible Virasoro module with given conformal dimension Δ_p . Let us consider at first the cases when p is positive integer or half-integer number. It is convenient to realize the irreducible Virasoro modules $\mathcal{L}_p, (\mathcal{L}_p^*)$ as a subspaces (factorspaces) $\text{Ker}_{\mathcal{F}_p}[X^l], \mathcal{F}_{-p}/\text{Im}[X^l]$ in the Fock modules where $l = p + \frac{1}{2}$ for R sector and $l = p + 1$ for NS sector. Then we have

Proposition 3.3

The following vertex operators of representation π_Z can be realized as the bosonic operators in the Fock submodules

i)

$$\begin{aligned}\Phi_{21}^+(\zeta) \mathbf{v} &= e^{i\phi(\zeta)} \mathbf{v}, \\ \Phi_{21}^{*+}(\zeta) \mathbf{v}^* &= \int_C \frac{dz}{2\pi iz} e^{-2i\phi(z)} e^{i\phi(\zeta)} \mathbf{v}^*,\end{aligned}\tag{3.50}$$

ii)

$$\begin{aligned}\Phi_{12}^{*-}(\zeta) \mathbf{v}^* &= e^{-\frac{i}{2}\phi(\zeta)} \mathbf{v}^*, \\ \Phi_{12}^-(\zeta) \mathbf{v} &= \int_C \frac{dz}{2\pi iz} e^{i\phi(z)} e^{-\frac{i}{2}\phi(\zeta)} \mathbf{v},\end{aligned}\tag{3.51}$$

here $\mathbf{v} \in \mathcal{L}_p$ and $\mathbf{v}^* \in \mathcal{L}_p^*$;

The integration contour C going counterclockwise is chosen to have the beginning and end at the origin of complex z -plane. It encloses all singularities of which positions are determined by the vector $\mathbf{v}^*(\mathbf{v})$. Notice, that due to this prescription the integrals are well defined.

This proposition is rather evident since part i) is a direct consequence of the equation (3.13) while the proof of part ii) is based on the following important property of operator $e^{-\frac{i}{2}\phi(\zeta)}$:

$$X|_{\pi_{NS}} e^{-\frac{i}{2}\phi(\zeta)} = -e^{-\frac{i}{2}\phi(\zeta)} X|_{\pi_R}.\tag{3.52}$$

Now we want to check our guess that formulae (3.50)-(3.51) describe the bosonization of chiral primaries for any $p > 0$. Let us note, that irreducible Virasoro algebra modules $\mathcal{L}_p$ ($\mathcal{L}_p^*$) in the case of generic p are isomorphic to Fock modules $\mathcal{F}_p$ ($\mathcal{F}_{-p}$). Indeed, the action of Virasoro algebra in $\mathcal{F}_p$ can be determined through the formulae (3.15) but there are no intertwining operators between Fock modules.³ Therefore operators (3.50)-(3.51) act from irreducible Virasoro algebra module into irreducible one. So, one need to check

³ For instance, the action of the integral operators like X, X' is not defined in these modules.

only the commutation relations of these operators with Virasoro algebra generators. But it are rather evident such as the integrals in (3.50)-(3.51) are well-defined in the general case too. Now straightforward computation prove that proposition 3.3 is still true for the chiral primary operators in the case of generic value of p .

Bosonization prescription (3.50)-(3.51) and conjugation condition (3.37) , (3.40) allow to work out the integral representation for any matrix element of operators $\Phi_{12}^{\pm}$ and $\Phi_{21}^{\pm}$. For instance, let us write down alternative derivation of the function $G_p'^{\pm}(\zeta)$. It would seems at first glance that calculation of matrix elements using the scalar product (3.29)-(3.30) is very non-trivial problem. The essence of the bosonization method is in the fundamental fact that the scalar product (3.21) restricted on the submodules of Fock modules

- i) $Ker_{\mathcal{F}_p}[X^l]$, $\mathcal{F}_{-p}/Im[X^l]$, $l = p + \frac{1}{2}$ for R sector, $l = p + 1$ for NS sector; p is positive integer or half-integer,
- ii) $\mathcal{F}_p$, $\mathcal{F}_{-p}$, for generic p ,

which are isomorphic to irreducible modules of Virasoro algebra, coincides with (3.30). Using this fact and (3.50)-(3.51) we get the formula:

$$(\mathbf{v}_p^*, \Phi_{12}^+(\zeta_2) \Phi_{12}^-(\zeta_1) \mathbf{v}_p) = \int_C \frac{dz}{2\pi i z} (\mathbf{f}_{-p}, e^{-\frac{i}{2}\phi(\zeta_2)} e^{i\phi(z)} e^{-\frac{i}{2}\phi(\zeta_1)} \mathbf{f}_p)_0 , \quad (3.53)$$

where the integration contour C is determined by the same prescription as in (3.51). The formula (3.53) leads to integral representation for functions $G_p'^+(\zeta)$ if we use the well-known rules for average of the exponent operators in the bosonic Fock space:

$$(\mathbf{f}_{-p}, e^{i\ell_n\phi(\zeta_n)} \dots e^{i\ell_1\phi(\zeta_1)} \mathbf{f}_p)_0 = \prod_{k>m} (\zeta_k - \zeta_m)^{\ell_k \ell_m} \prod_k \zeta_k^{\frac{\ell_k^2}{2} + \ell_k p} \delta_{\ell_1 + \dots + \ell_n, 0} . \quad (3.54)$$

Then the function (3.53) can be rewritten in term of hypergeometric function via integral representation:

$$\int_C \frac{dz}{2\pi i} z^{c-1} (1-z^{-1})^{-a} (1-\zeta z)^{-b} = \frac{\Gamma(c+a)}{\Gamma(c+1)\Gamma(a)} F(a+c, b, c+1; \zeta) \quad Re[c] > 0. \quad (3.55)$$

To find the function $G_p'^-$ we can use the relation

$$[G_p'^+(\zeta^{-1})]^* = G_{-p}'^-(\zeta), \quad |\zeta| = 1 , \quad (3.56)$$

which follows from the conjugation conditions (3.40). In this way we reproduce formula (3.45).




Fig. 3: Integration contour for hypergeometric function.

4. Deformed Virasoro algebra with $c=-2$.

In the previous section we have developed bosonization procedure to study the representations of ZF algebra. The worth important step there was the introduction of integral operators X, X' acting in the bosonic space $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{\frac{k}{2}}$. In this space we introduced the action of $c = -2$ Virasoro algebra by defining generators $T(z) = \sum L_n z^{-n-2}$ as composite operators built from Heisenberg generators b_n . The property, that Virasoro algebra generators commute with X and X' allowed us classify the representation space of ZF algebra by the action of *Symm* algebra. Moreover, we pointed out that operators $Z_{\pm}$ itself can be constructed as some linear combinations of vertex operators Φ_{21} of Virasoro algebra. We also determined another elementary chiral primary Φ_{12} the commutation relations of which are defined by the constant R-matrix of IRF type.

Of course, representations of algebra (3.1) - (3.2) could be investigated directly in the fermionic language. However, the bosonization procedure can be applied in constructions where fermionic description is lacking. One of the lessons we have studied above is that ZF algebra, Virasoro algebra, algebras of chiral primary operators and algebras of screening operators are mutually related in the bosonic picture. We would like to examine further these interrelations considering less trivial situations.

It will be convenient to start with bosonic space and deformed integral operators X and X' rather than with ZF algebra. Indeed, operators X and X' could be treated as completely specifying the construction. For instance, universal enveloping algebra of Virasoro algebra could be considered as subalgebra of operators from Heisenberg algebra commuting with X and X' , while operators Φ_{21}, Φ_{12} were specified with respect to Virasoro algebra, etc. From this point of view, generalizations of the proposed construction are determined by the appropriate deformations of screening operators.

The well-known generalization of proposed bosonic construction is originated in deformation of operators X and X' in a way which leads to Virasoro algebra with central charge $-2 < c < 1$. This direction will be discussed later.

There exists, however, another class of deformations [17] relating with generalizations of the Virasoro algebra. These deformations will be called hereafter by x-deformations. Again, x-deformation is determined by the appropriate continuous deformation of screening operators. In distinction with Virasoro algebra cases, x-deformation leads to more general associative algebras of deformed vertex operators Φ_{21}, Φ_{12} .

In the present section we want to elaborate simplest example of x-deformation corresponding to $c = -2$ Virasoro algebra. In this particular instance the deformation has remarkably simple form. Working out this toy model we try to extract information on essential properties of general x-deformation. Namely, we argue that the whole Fock space can be decomposed under the action of operators X and X' into direct sum of irreducible representations of deformed Virasoro algebra proposed in [30]. This algebra is determined as subalgebra of operators from the universal enveloping algebra of Heisenberg algebra which commute with X and X' . It should be clear that such commutants always generate associative algebra. Indeed, the linear space of commuting with X and X' operators is closed under the operation of multiplication, while associativity condition follows from the fact that $T(\zeta)$ are build upon generators of the Heisenberg algebra. Hence, in general case one can also expect that Fock space will be divided into direct sum of representations of some infinite-dimensional algebra generalizing Virasoro algebra.

Another purpose of this example is to demonstrate how one can find the deformation of chiral primary operators Φ_{12} and Φ_{21} . More explicitly, how to obtain bosonic realization for such operators. We will show that commutation relations of operators Φ_{12} are determined by elliptic R-matrix of IRF type. This justify proposed deformation.

Further we will use the same notations as an earlier, pointing only those definitions and formulae which contain significant distinctions. For instance, symbol $\mathcal{L}$ will mean the irreducible representation of deformed $c = -2$ Virasoro algebra, Φ_{12} will be deformation of correspondent chiral primary etc.

4.1. Deformation of the Virasoro algebra with $c = -2$

Let us start with consideration of the bosonic space $\pi_Z = \oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}$ occurred in the section 3. The x -deformation of the construction given in the previous section is determined by the following redefinition of the operator X :

$$X = \oint \frac{dz}{2\pi iz} e^{-i(\phi(zx) + \phi(zx^{-1}))} . \quad (4.1)$$

where deformation parameter is a real number $0 < x < 1$. If second operator X' is still unchanged, then it is not hard to see that formulae (3.10) - (3.12) are valid. Let us fix notations of new operators, exact sense of which will become transparent later:

$$\begin{aligned} Z_+(\zeta) &= e^{i\phi(\zeta)} , \\ Z_-(\zeta) &= [X, Z_+(\zeta)] \end{aligned} \quad (4.2)$$

The integral in the definition of Z_- could be computed explicitly:

$$Z_-(\zeta) = \frac{e^{-i\phi(\zeta x^2)} - e^{-i\phi(\zeta x^{-2})}}{x - x^{-1}} . \quad (4.3)$$

In accordance with the scheme of previous section let us describe at first the structure of the subspaces $\text{Ker}_{\mathcal{F}_{l-\frac{1}{2}}^+} [X^l]$ in Fock space. To do this one need find operators which would generalize (3.14). It might be easily checked that the action of the following operators in the Fock space π_Z commute with X and X' :

$$-\zeta^2 T^\nu(\zeta) = \frac{Z_+(\zeta x^{-2\nu-1})Z_-(\zeta x^{2\nu+1}) - Z_-(\zeta x^{-2\nu-1})Z_+(\zeta x^{2\nu+1})}{2(x + x^{-1})} , \quad (4.4)$$

where $\nu \in \mathbf{Z}$. Note that

$$T^\nu(\zeta) \equiv T^{-\nu-1}(\zeta) .$$

In the limit $x \rightarrow 1$ operators $T^\nu(\zeta)$ become Virasoro algebra generators expressed by formula (3.15). Straightforward calculation ensure that modes of the Laurent expansion $T^\nu(\zeta) = \sum_{n \in \mathbf{Z}} L_n^\nu \zeta^{-n-2}$ generate so-called deformed Virasoro algebra [30]:

$$\begin{aligned} 2[L_n^\nu, L_m^\mu] &= [(\mu + 1)n - (\nu + 1)m]_{x^2} L_{n+m}^{\nu+\mu+1} + [(\mu + 1)n + \nu m]_{x^2} L_{m+n}^{\mu-\nu} - \\ &- [\mu n + (\nu + 1)m]_{x^2} L_{n+m}^{\nu-\mu} - [\mu n - \nu m]_{x^2} L_{n+m}^{\nu+\mu} - \\ &- \frac{\delta_{m+n,0}}{4(x^2 - x^{-2})^2} (C_n^{\nu,\mu} + C_n^{\nu,-1-\mu} + C_n^{-1-\nu,\mu} + C_n^{-1-\nu,-1-\mu}) , \end{aligned} \quad (4.5)$$

where

$$C_n^{\nu,\mu} = \frac{[n(\mu + \nu)]_{x^2}}{[\mu + \nu]_{x^2}} - \frac{[n(\nu - \mu)]_{x^2}}{[\nu - \mu]_{x^2}}.$$

We denote this algebra as $Vir_{-2,x}$. Let us adjoin to the algebra (4.5) a derivation D with property $[D, L_n^\nu] = nL_n^\nu$ for any $\nu \in \mathbf{Z}$. In the bosonic Fock spaces this operator can be realized as $D = \sum_{m>0} b_{-m}b_m + \frac{P^2}{2} - \frac{1}{8}$. It provides the universal enveloping algebra of $Vir_{-2,x}$ with structure of $\mathbf{Z}$ -graded algebra. Then the triangular decomposition of $Vir_{-2,x}$ is a decomposition into elements of positive ($L_n^\nu, n < 0$), zero (D, L_0^ν) and negative ($L_n^\nu, n > 0$) degrees for any $\nu \in \mathbf{Z}$. By the definition, Verma module of the deformed Virasoro algebra is a $\mathbf{Z}$ -graded module generated by operators $L_n^\nu, n < 0$ under the action on the unique highest weight vector $\mathbf{v}$. Verma module inherits structure of $\mathbf{Z}$ -graded space from the grading of universal enveloping algebra. Vector $\mathbf{v}$ is the highest weight vector of Verma module with weight Δ if it is annihilated by any operator $L_n^\nu, n > 0$, and the action of elements with zero grading on it, yields $L_0^\nu \mathbf{v} = \Delta^\nu \mathbf{v}$; $D\mathbf{v} = \Delta \mathbf{v}$. In general situation Verma module is degenerated, i.e. it contains invariant subspaces created by action of operators $L_n^\nu, n < 0$ on null vectors. Null vectors, by definition, obey the equation $L_n^\nu \mathbf{v}^0 = 0, n > 0$; $D\mathbf{v}^0 = (\Delta + N)\mathbf{v}^0$ with some integer N (degeneration level). In order to obtain an irreducible representation, we have to put all null-vectors together with the whole subspace generated by it equal to zero; that is we have to factorize Verma module by all invariant submodules.

In spite of the fact that operator X and its commutants in the Fock space have been deformed, the description of structure of Fock space $\oplus_k \mathcal{F}_{k-1/2}^+$ in terms of irreducible $Vir_{-2,x}$ modules is almost the same as an earlier. Indeed, let generators L_n^ν are realized through (4.4). Then one can demonstrate that the highest vectors $\mathbf{f}_{l-\frac{1}{2}}$ of Fock module turn to be highest vectors of the Verma modules of $Vir_{-2,x}$. It is possible to show that acting on the highest vectors of Fock space $\mathbf{f}_{l-\frac{1}{2}} \in \mathcal{F}_{l-\frac{1}{2}}^+, l = 1, 2, \dots$ generators L_n^ν create subspaces $Ker_{\mathcal{F}_{l-\frac{1}{2}}^+} [X^l]$ which are isomorphic to irreducible $Vir_{-2,x}$ modules $\mathcal{L}_{l-\frac{1}{2}}$ correspondingly. Notice, that the character of irreducible module of $Vir_{-2,x}$ coincides with the character of $c = -2$ Virasoro algebra:

$$Tr_{\mathcal{L}_{l-\frac{1}{2}}} [q^{D-\frac{c}{24}}] = q^{\frac{1}{12}} \frac{q^{\frac{l(l-1)}{2}} - q^{\frac{l(l+1)}{2}}}{\prod_{n=1}^{\infty} (1 - q^n)}. \quad (4.6)$$

The numbers of the states on given level will be conserved in a general x -deformation too. Recall, that similar situation take place in the representation theory of quantum groups

[26], [27], [28] where characters of deformed irreducible modules remain to be the same as undeformed one if the deformation parameter is not a primitive root of unit.

The spaces $\mathcal{F}_{l-\frac{1}{2}}^+$ and $\mathcal{F}_{l+1/2}^*$ can be decomposed into the direct sum of $Vir_{-2,x}$ modules analogously to (3.23) and (3.20) where, of course, we should bear in the mind that symbol $\mathcal{L}_{l-\frac{1}{2}}$ means now irreducible module of deformed Virasoro algebra.

Now the whole algebra $Symm$ acting on the space $\pi_Z^R \oplus \pi_Z^{R*}$ is the tensor product of infinite-dimensional deformed Virasoro algebra with $c = -2$, $0 < x < 1$ and finite-dimensional part generated by operators X, X^*, X', X'^* and P with commutation relations (3.24) - (3.27). For this reason, the decomposition of the space $\pi_Z^R \oplus \pi_Z^{R*}$ into direct sum of irreducible representation of algebra $Symm$ has the form (3.28). The scalar product in the space $\oplus_{k \in \mathbf{Z}} \mathcal{F}_{k-\frac{1}{2}}$ is given by (3.29) and the conjugation conditions $(\mathbf{v}_1 L_n^\nu, \mathbf{v}_2) = (\mathbf{v}_1, L_{-n}^\nu \mathbf{v}_2)$. Moreover, in the calculations one can use the scalar product (3.21). The arguments here are practically the same as in subsec.3.6.

In the NS sector the finite-dimensional part of $Symm$ algebra is still given by the $sl(2)$ algebra and similarly to undeformed situation we have isomorphism $Ker_{\mathcal{F}_l} X^l \cong \mathcal{L}_l$. In addition, the decomposition of Fock spaces into direct sum of irreducible modules of deformed Virasoro algebra is described by the formula (3.33) again. The scalar product in the space π_Z^{NS} is the same as (3.29)-(3.30) and the space π_Z^{NS} can be treated as selfdual $\pi_Z^{NS} = \pi_Z^{NS*}$.

4.2. Deformation of vertex operators

Now we want to establish x -deformation of chiral primary operators $\Phi_{21}^\pm$ and $\Phi_{12}^\pm$. The construction of first operator $\Phi_{21}^\pm$ is rather evident. Indeed, it can be realized in the Fock spaces in the same manner as (3.50):

$$\begin{aligned} \Phi_{21}^+(\zeta) \mathbf{v} &= e^{i\phi(\zeta)} \mathbf{v}, \quad \mathbf{v} \in \mathcal{L}_p, \\ \Phi_{21}^{*+}(\zeta x^2) \mathbf{v}^* &= \int_C \frac{dz}{2\pi i z} e^{-i[\phi(zx) + \phi(zx^{-1})]} e^{i\phi(\zeta)} \mathbf{v}^*, \quad \mathbf{v}^* \in \mathcal{L}_p^*, \end{aligned} \quad (4.7)$$

where we explicitly show the states on which such operators are well-defined. As an earlier, we take contours in the integral $\Phi_{21}^{*+}(\zeta)$ with beginning and ending in zero, and enclose all singularities which positions are determined by vector v^* . Demanding conjugation properties of $\Phi_{21}^\pm(\zeta)$ for $|\zeta| = 1$ as

$$(\mathbf{u}^* \Phi_{21}^{*\pm}(\zeta x^2), \mathbf{v}) = (\mathbf{u}^*, \Phi_{21}^\mp(\zeta) \mathbf{v}), \quad (4.8)$$

where $\mathbf{u}^* \in \mathcal{L}_p^*$ and $\mathbf{v} \in \mathcal{L}_{p\pm 1}$, one can easily obtain any matrix element of such operators. The bosonic realization (4.7) possesses to find that commutation relations of these vertex operators with deformed Virasoro algebra are given by formula

$$2 \zeta^{-n} [L_n^\nu, \Phi_{21}^\pm] = \frac{x^{2\nu+3} \Phi_{21}^\pm(\zeta x^{4\nu+4}) - x^{-2\nu-3} \Phi_{21}^\pm(\zeta x^{-4\nu-4})}{x^2 - x^{-2}} - \frac{x^{2\nu-1} \Phi_{21}^\pm(\zeta x^{4\nu}) - x^{-2\nu+1} \Phi_{21}^\pm(\zeta x^{-4\nu})}{x^2 - x^{-2}}. \quad (4.9)$$

To define deformed chiral primary operators Φ_{12} (3.39), let us remind that the crucial property in the proving of the proposition 3.3 was the equation (3.52). We demand that this basic relation is preserved in the deformed theory in the following sense. The second chiral primary have to be constructed in terms of new field ϕ' such that $\lim_{x \rightarrow 1} \phi'(z) = -\frac{1}{2}\phi(z)$ and

$$X|_{\pi_{NS}} e^{i\phi'(\zeta)} = -e^{i\phi'(\zeta)} X|_{\pi_R}. \quad (4.10)$$

Let ϕ' is built from generators P, Q, b'_n as:

$$\phi'(\zeta) = -\frac{1}{2}(Q - i P \ln \zeta) - \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} \frac{b'_m}{i m} \zeta^{-m}, \quad (4.11)$$

then condition (4.10) is obviously satisfied if new creation-annihilation operators be

$$b'_n = (x^n + x^{-n})^{-1} b_n. \quad (4.12)$$

Notice, that, due to this definition, the formula expressing X' in terms of field $\phi'(\zeta)$ has the form:

$$X' = \oint \frac{dz}{2\pi i z} e^{-i [\phi'(zx) + \phi'(zx^{-1})]}. \quad (4.13)$$

Comparing this with equation (4.1) one can observe that integral operators X, X' are built from ϕ and ϕ' correspondingly in the remarkably symmetrical form. Further we will see that this property allows direct generalization for the arbitrary x -deformation with $0 < x < 1$. Now the chiral primary operators, intertwining R and NS sectors can be realized as

$$\begin{aligned} \Phi_{12}^{*-}(x^2 \zeta) \mathbf{v}^* &= e^{i\phi'(\zeta)} \mathbf{v}^*, \quad \mathbf{v}^* \in \mathcal{L}_p^*, \\ \Phi_{12}^-(\zeta) \mathbf{v} &= \int_C \frac{dz}{2\pi i z} e^{-i [\phi'(zx) + \phi'(zx^{-1})]} e^{i\phi'(\zeta)} \mathbf{v}, \quad \mathbf{v} \in \mathcal{L}_p. \end{aligned} \quad (4.14)$$

The integral in $\Phi_{12}^-(\zeta)$ would be well-defined if contour is chosen as in (3.51). These bosonic prescriptions together with conjugation condition

$$(\mathbf{u}^* \Phi_{12}^{\star \pm}(\zeta x^2), \mathbf{v}) = (\mathbf{u}^*, \Phi_{12}^{\mp}(\zeta) \mathbf{v}) , \quad (4.15)$$

where $|\zeta| = 1$, completely fix the action of the operators $\Phi_{12}^{\pm}(\zeta)$ in the Fock space. The matrix elements of products of such operators can be derived by the standard bosonization technique. We leave the explicit calculations till the section 6 where it will be worked out as particular $\xi = 1$ case in context of general x -deformation. So, let us just note, that knowing matrix elements one can obtain by standard way the commutation relations of these operators. The essential difference of this case in comparison with undeformed one is that these commutation relations are determined by elliptic R-matrix of the IRF type rather than constant R-matrix:

$$\Phi_{12}^a(\zeta_1) \Phi_{12}^b(\zeta_2)|_{\mathcal{L}_p} = \sum_{c+d=a+b} \mathbf{W}' \left[\begin{matrix} p + \frac{a+b}{2} & p + \frac{c}{2} \\ p + \frac{b}{2} & p \end{matrix} \middle| \frac{\zeta_1}{\zeta_2} \right] \Phi_{12}^d(\zeta_2) \Phi_{12}^c(\zeta_1)|_{\mathcal{L}_p} . \quad (4.16)$$

The nontrivial elements of the matrix $\mathbf{W}'$ read:

$$\begin{aligned} \mathbf{W}' \left[\begin{matrix} p \mp 1 & p \mp \frac{1}{2} \\ p \mp \frac{1}{2} & p \end{matrix} \middle| \zeta \right] &= r(\zeta) , \\ \mathbf{W}' \left[\begin{matrix} p & p \mp \frac{1}{2} \\ p \mp \frac{1}{2} & p \end{matrix} \middle| \zeta \right] &= r(\zeta) \zeta^{\pm p - \frac{1}{2}} \frac{\Theta_{x^4}(x^2) \Theta_{x^4}(x^{\pm 4p} \zeta)}{\Theta_{x^4}(x^2 \zeta) \Theta_{x^4}(x^{\pm 4p})} , \\ \mathbf{W}' \left[\begin{matrix} p & p \pm \frac{1}{2} \\ p \mp \frac{1}{2} & p \end{matrix} \middle| \zeta \right] &= -r(\zeta) x^{\pm 2p} \zeta^{-\frac{1}{2}} \frac{\Theta_{x^4}(x^{2(\pm 2p+1)}) \Theta_{x^4}(\zeta)}{\Theta_{x^4}(x^{\pm 4p}) \Theta_{x^4}(x^2 \zeta)} , \end{aligned} \quad (4.17)$$

where

$$r(\zeta) = \zeta^{\frac{1}{4}} \exp \left[\sum_{m=1}^{+\infty} \frac{\zeta^m - \zeta^{-m}}{m (x^m + x^{-m})^2} \right] .$$

The associativity of algebra (4.16) follows from the fact that matrix $\mathbf{W}'$ is a solution of Yang-Baxter equation. This justify proposed deformation of chiral primary operators Φ_{12} .

4.3. The x -deformation of fermions ZF algebra

So far we were interested in vertex operators of $Vir_{-2,x}$ algebra. To construct vertex operators of $Symm$ algebra, one need take into account remaining finite-dimensional part of symmetry algebra.

Let $\mathbf{e}_l^m$ be basic vectors of irreducible spin $j = \frac{l-1}{2}$ representation $\mathcal{V}_l$ of algebra $sl(2)$ and $\mathbf{v} \in \mathcal{L}_l$. Define operators $Z_{\pm}(\zeta)$ acting on the space π_Z^R and π_Z^{NS} by the formula

(3.35). The straightforward computation shows that operators $Z_{\pm}(\zeta)$ are generators of ZF algebra given by commutation relations

$$Z_a(\zeta_1)Z_b(\zeta_2) = -Z_b(\zeta_2)Z_a(\zeta_1), \quad \zeta_1 \neq \zeta_2 \quad (4.18)$$

and operator product expansion:

$$\begin{aligned} Z_{\pm}(\zeta_2)Z_{\mp}(\zeta_1) &= \pm \frac{(x + x^{-1}) \zeta_1 \zeta_2}{(\zeta_2 - \zeta_1 x^2)(\zeta_2 - \zeta_1 x^{-2})} + O(1), \\ Z_{\pm}(\zeta_2)Z_{\pm}(\zeta_1) &= O(1). \end{aligned} \quad (4.19)$$

It is rather evident now that operators (4.2) introduced in the beginning of this section are exactly generators of ZF algebra (4.18)-(4.19).

5. Nondeformed Virasoro algebra with $c < 1$

In the present section we remind how to construct representations of ZF algebra (more explicitly, a pair of ZF algebras) with constant R matrix corresponding to quantum group $U_q(sl(2))$. As an earlier, the representation space of ZF algebra can be realized in the direct sum of irreducible representations of Virasoro algebra with central charge $c < 1$. The operators of ZF algebras will be expressed in terms of vertex operators of Virasoro algebra Vir_c . Therefore, the main objects of our investigation will be the irreducible representations of Virasoro algebra and vertex operator algebra. To describe these objects we will use bosonization method, examples of which was presented in the previous sections.

In the case under the consideration the explicit realization of Virasoro algebra generators L_n in the bosonic Fock space $\mathcal{F}_p$ is well-known [2]. Then universal enveloping algebra $U(Vir_c)$ turn to be subalgebra in the universal enveloping algebra of Heisenberg algebra. Moreover, any highest weight vector in the Fock module will be the highest weight vector of Verma module of Vir_c . The original motivation of bosonization of Virasoro algebra [9] is that in the Fock space one can explicitly construct the intertwining operators between Verma modules. In the physical literature these operators were historically called by screening operators [2]. According to the definition, intertwining operators commute with any element of $U(Vir_c)$, hence it have to map singular vectors of Verma module into singular ones. So the analysis of the structure of the reducible Verma modules becomes very simple. Indeed, to determine singular vectors one needs to find such vectors in the Fock space which are mapped under the action of screening operators into highest

weight vectors, or such vectors which can be obtained from the highest weight vectors by the action of screening operators. For this reason, knowledge of all possible intertwining operators is equivalent to the knowledge of the invariant subspaces in the Verma modules. This subject was intensively discussed in the literature. It is well known, that structure of Verma modules of Vir_c drastically depends on the arithmetical properties of real number c . Having in the mind further x -deformation we will omit complicated cases of completely reducible Verma modules just considering the generic case, i.e. case 2 in the Feigin-Fuks classification [9]. Our task is to remind the essential features which occur in the bosonization of representations of Virasoro algebra and ZF algebra. (The analysis of more complicated cases can be found in the refs. [16], [31]). In particular we would like to clear up the idea that knowledge of the intertwining operators between Verma modules of Vir_c is sufficient to describe the irreducible representations of Virasoro algebra and algebra of vertex operators without appealing to the commutation relations of generators L_n . Another very important point we recall and constantly use in the construction is the existence of remarkable discrete symmetry which in our notations is just a change $\xi \leftrightarrow -1 - \xi$ ($\alpha_- \leftrightarrow \alpha_+$ in the Dotsenko-Fateev work [2]). This symmetry, origin of which is still hardly understood, seems to be essential not only in CFT [29] but also in general theory of integrable models [17], [18].

Our present consideration is based on slightly different point of view on the bosonization (see also section 3,4). Namely, we consider screening operators as basic objects which completely determine the whole construction. In particular, Virasoro algebra can be treated as an algebra of generators in the Fock space which commute with correspondent screening operators. In this approach we can investigate the irreducible representations of Virasoro algebra independently on the facts what basis in $U(Vir_c)$ or what commutation relations are really involved. The similar idea was applied in the development of the theory of W-algebra [32]. So, the most important step is the introduction of correctly defined screening operators. These operators have remarkably simple form. They are given by some powers of operators X and X' which are deformations of (3.9). We explicitly enlist the subspaces in the Fock space where these integral operators are well-defined. As soon as we determine screening operators, we are able to describe the irreducible representations of Virasoro algebra as some submodules (or factor modules) of Fock modules. Our next step is the definition of the chiral vertex operators of Virasoro algebra. To do this, we need fix basis L_n in the $U(Vir_c)$ since, in general, the knowledge of intertwining operators is not

enough to uniquely determine vertex operators. Our task is to extract necessary properties of these operators which are determined by screening operators rather than by choice of basis L_n . Defining the bosonization of chiral primaries, we demonstrate that matrix elements of these operators can be easily computed using Wick theorem. The studying the analytical properties of four point functions shows that commutation relations of chiral primary operators Φ_{12} , Φ_{21} are determined by two different constant $\mathbf{W}$ -matrices which are solutions of Yang-Baxter equation of IRF type. To construct the ZF algebras of "vertex type" [8] one needs take into account multiplicities of irreducible representations of Virasoro algebra. We argue that there are two ZF algebras associated to the algebras of chiral primaries Φ_{12} and Φ_{21} correspondingly. The R-matrices of these ZF algebras correspond to the R-matrices of quantum groups $U_p(sl(2))$ with different deformation parameters p [29]. These R-matrices are connected by the transformation $\xi \leftrightarrow -1 - \xi$ which is originated from the symmetry between Φ_{12} and Φ_{21} (or $\phi' \leftrightarrow \phi$). The irreducible representation of a pair of ZF algebras coincides with direct sum of Fock modules, hence it admit classification by the representations of the Virasoro algebra.

The bosonization is just a useful method to study the representations of Virasoro and ZF algebras. Of course, the results do not depend on it. In particular, without any bosonization the irreducible representations of a pair of $[\xi \leftrightarrow -1 - \xi]$ -symmetrical ZF algebras are isomorphic to direct sum of irreducible representations of Virasoro algebra.

This construction can be considered as case $x = 1$ of general two parametric deformation with parameters x and ξ . In the following section we will generalize main statements for $x \neq 1$ deformation.

3.1 Let us introduce the free bosonic field

$$\phi(z) = \sqrt{\frac{\xi + 1}{2\xi}} (Q - i P \ln z) + \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} \frac{b_m}{i m} z^{-m}, \quad (5.1)$$

where the commutation relation of the zero modes P, Q is defined by $[Q, P] = i$, while determining relation for modes b_n is:

$$[b_m, b_n] = \frac{\xi + 1}{2\xi} m \delta_{m+n, 0}. \quad (5.2)$$

Further we will consider cases with deformation parameter $\xi > 1$. Moreover, in order to avoid additional complications we assume that this number is irrational. Note that the

case $\xi = 1$ corresponds to the example discussed in the section 3. Together with the field ϕ , it is convenient to define another one:

$$\phi'(z) = -\sqrt{\frac{\xi}{2(\xi+1)}} (Q - i P \ln z) - \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} \frac{b'_m}{i m} z^{-m}, \quad (5.3)$$

where

$$\xi b_m = (\xi + 1) b'_m.$$

As we will see below, there is a remarkable symmetry with respect the transformation $\xi \leftrightarrow -1 - \xi$. This is the reason why we introduce independent notation for field $\phi'(z)$. In the case $x = 1$ fields $\phi'(z)$ and $\phi(z)$ are just proportional. However, in the x-deformed construction their connection turns to be more complicate.

In the set of Fock modules

$$\mathcal{F}_{k,k'} \equiv \mathcal{F}_{\frac{(\xi+1)k-\xi k'}{\sqrt{2\xi(\xi+1)}}}, \quad k, k' \in \mathbf{Z} \quad (5.4)$$

one might introduce the following formal operators:

$$\begin{aligned} X^l &= \int_{C_1} \dots \int_{C_l} \frac{dz_1}{2\pi i z_1} \dots \frac{dz_l}{2\pi i z_l} e^{-2i\phi(z_1)} \dots e^{-2i\phi(z_l)}, \\ X'^{l'} &= \int_{C_1} \dots \int_{C_{l'}} \frac{dz_1}{2\pi i z_1} \dots \frac{dz_{l'}}{2\pi i z_{l'}} e^{-2i\phi'(z_1)} \dots e^{-2i\phi'(z_{l'})}. \end{aligned} \quad (5.5)$$

Here we used Felder's prescription [16] for integration contours (see fig.4.). Namely, any contour C_i , $1 \leq i \leq l-1$ has beginning and ending in the point z_l chosen on the unit circle. It encloses the origin and all singularities depending on the variables $z_k (k < i)$. The last integration on the variable z_l is provided along the circle C_l .

In essential distungish with the case $c = -2$ the action of these operators is ill defined in the whole set of Fock modules since the last integration contours $C_l (C_{l'})$ in the definitions (5.5) is not closed. Hence, first of all, we must formulate where the operators (5.5) act.

Proposition 5.1

The action of operator $X^l (X'^{l'})$ is defined only on the Fock modules $\mathcal{F}_{l,k} (\mathcal{F}_{k,l'}), k \in \mathbf{Z}$ and $l, l' > 0$. Then

$$\begin{aligned} Ker_{\mathcal{F}_{l,l'}}[X^l] &= Ker_{\mathcal{F}_{l,l'}}[X'^{l'}], \\ Ker_{\mathcal{F}_{l,-l'}}[X^l] &= Ker_{\mathcal{F}_{l,-l'}}[X'^{l'}] = 0, \\ Im_{\mathcal{F}_{l,l'}}[X^l] &= \mathcal{F}_{-l,l'}, \\ Im_{\mathcal{F}_{l,l'}}[X'^{l'}] &= \mathcal{F}_{l,-l'}, \\ Im_{\mathcal{F}_{l,-l'}}[X^l] &= Im_{\mathcal{F}_{l,-l'}}[X'^{l'}] \subset \mathcal{F}_{-l,-l'}. \end{aligned} \quad (5.6)$$




Fig. 4: The integration contours.

We will call integral operators (5.5) defined on the correspondent spaces by screening operators. Considering the screening operators as basic objects, one can define Virasoro algebra as follows:

Definition. Let operators $P, Q, b_n, n \in \mathbf{Z}$ satisfy the commutation relations (5.2) and screening operators are given by (5.5). Virasoro algebra is a subalgebra in the universal enveloping algebra of Heisenberg algebra of operators b_n and P . An element from universal enveloping of Heisenberg algebra belongs to Virasoro algebra if it commutes with an action of the screening operators.

In present case one can simply write down explicit expressions for basic generators of the space of invariants in the universal enveloping of Heisenberg algebra:

$$\begin{aligned}
 L_n &= \sum_{k+m=n} b_m b'_k - (\sqrt{2\xi(\xi+1)}P + n)(b_n - b'_n), \quad n \neq 0; \\
 L_0 &= \sum_{m>0} (b_{-m} b'_m + b'_{-m} b_m) + \frac{P^2}{2} - \frac{1}{4\xi(\xi+1)}.
 \end{aligned}
 \tag{5.7}$$

The L_n obey Virasoro commutation relations with the central charge $c = 1 - \frac{6}{\xi(\xi+1)}$. So, the Fock space is given the structure of Virasoro module. The highest Fock state $\mathbf{f}_{k,k'} \equiv \mathbf{f}_{\frac{(\xi+1)k - \xi k'}{\sqrt{2\xi(\xi+1)}}}$ will also be the highest weight vector of Verma module of Vir with the conformal dimension

$$\Delta_{k,k'} = \frac{((\xi+1)k - \xi k')^2 - 1}{4\xi(\xi+1)}.
 \tag{5.8}$$

The most important property of operators (5.5) is that they commute with generators of the form (5.7) , i.e. are intertwining operators between Verma modules of the Virasoro algebra. Knowing intertwining operators one might study the structure of Verma modules $\mathcal{M}_{k,k'}$ of Vir_c and construct the irreducible representations $\mathcal{L}_{k,k'}$ as some subspaces or factor spaces of Fock spaces. Namely, introduce the following notations:

$$\mathcal{L}_{k,k'} = \begin{cases} Ker_{\mathcal{F}_{k,k'}}[X^k], & \text{if } k, k' \geq 1; \\ \mathcal{F}_{k,k'} / Im_{\mathcal{F}_{-k,k'}}[X^{-k}], & \text{if } k, k' \leq -1; \\ \mathcal{F}_{k,k'}, & \text{otherwise,} \end{cases} \quad (5.9)$$

We claim that the following proposition is hold:

Proposition 5.2

Let generators of Virasoro algebra are given by formulae (5.7) and parameter $\xi > 1$ be irrational number. Then

1. *The space $\mathcal{L}_{k,k'}$ for any integer number k, k' is irreducible representation of Virasoro algebra with central charge $c = 1 - \frac{6}{\xi(\xi+1)}$. The highest weight vector of $\mathcal{L}_{k,k'}$ coincides with the vector $\mathbf{f}_{k,k'}$ and has the conformal dimensions (5.8).*
2. *For $k \cdot k' \leq 0$ or $k, k' < 0$, the Verma module $\mathcal{M}_{k,k'}$ built upon highest weight vector $\mathbf{f}_{k,k'}$ coincides with the total Fock module $\mathcal{F}_{k,k'}$.*

We would like now to comment propositions 5.1, 5.2 [31]. As we have noted, any highest weight vector $\mathbf{f}_{k,k'}$ of Fock module $\mathcal{F}_{k,k'}$ turns to be highest weight vector of Verma module $\mathcal{M}_{k,k'}$ of Virasoro algebra. Generically speaking, Fock space $\mathcal{F}_{k,k'}$ does not coincide neither with Verma module $\mathcal{M}_{k,k'}$, nor with irreducible module $\mathcal{L}_{k,k'}$ of Vir_c . It might contain some subspaces which are invariant with respect to the action of generators (5.7). Such as screening operators (5.5) commute with any generator from universal enveloping of Vir_c then they have to map invariant subspace of the Vir into invariant one. In the case under the consideration the structure of embedding of Verma modules given by the actions of screening operators is rather simple:

- (i) Consider at first Fock module $\mathcal{F}_{l,l'}$, $l, l' > 0$. According to proposition 5.1 this module contains a vector $\mathbf{f}_{l,l'}^0$ such that $X^l \mathbf{f}_{l,l'}^0 = \mathbf{f}_{-l,l'}$ ($X'^{l'} \mathbf{f}_{l,l'}^0 = \mathbf{f}_{l,-l'}$). This vector can not be obtained by the action of the Virasoro algebra generators on the highest weight vector $\mathbf{f}_{l,l'}$ and Fock submodule built upon state $\mathbf{f}_{l,l'}^0$ turns to be invariant space with respect to the action of generators of Virasoro algebra. Indeed, if $\mathbf{f}_{l,l'}^0$ is produced by the action of any element from universal enveloping of Vir_c on the vector $\mathbf{f}_{l,l'}$, then operator X^l ($X'^{l'}$) would map it into zero rather than into $\mathbf{f}_{-l,l'}$ ($\mathbf{f}_{l,-l'}$), since X^l ($X'^{l'}$) commutes with any

generator of Vir_c and $X^l \mathbf{f}_{l,l'} = X'^{l'} \mathbf{f}_{l,l'} = 0$. The proposition 5.2 means that subspace $Ker_{\mathcal{F}_{l,l'}}[X^l] = Ker_{\mathcal{F}_{l,l'}}[X'^{l'}]$ is irreducible representation $\mathcal{L}_{l,l'}$ of Virasoro algebra. The modules $\mathcal{F}_{-l,l'} = Im_{\mathcal{F}_{l,l'}} X^l$ and $\mathcal{F}_{l,-l'} = Im_{\mathcal{F}_{l,l'}} X'^{l'}$ coincide with Verma modules of Vir_c and do not have any invariant subspaces. Therefore it can be identified with $\mathcal{L}_{l,-l'}$ and $\mathcal{L}_{-l,l'}$ correspondingly.

(ii) The Fock space $\mathcal{F}_{-l,-l'}$ is a reducible Verma module of Vir_c with unique null-vector $\mathbf{f}_{-l,-l'}^0$ which is the image of the highest weight vector $\mathbf{f}_{l,-l'}$ ($\mathbf{f}_{l,-l'}$) under the action of the operator X^l ($X'^{l'}$):

$$\mathbf{f}_{-l,-l'}^0 = X^l \mathbf{f}_{l,-l'} = X'^{l'} \mathbf{f}_{-l,l'} \in \mathcal{F}_{-l,-l'}.$$

The irreducible representation $\mathcal{L}_{-l,-l'}$ of Vir turn to be isomorphic to factor space $\mathcal{F}_{-l,-l'} / Im_{\mathcal{F}_{l,-l'}}[X^l] = \mathcal{F}_{-l,-l'} / Im_{\mathcal{F}_{-l,l'}}[X'^{l'}]$ of Fock space.




Fig. 5: The structure of spaces $\mathcal{F}_{l,l'}$ and $\mathcal{F}_{-l,-l'}$

The most important objects for us are the sets of irreducible representations $\{\mathcal{L}_{l,l'} | l, l' > 0\}$ and $\{\mathcal{L}_{l,l'}^* | l, l' > 0\}$ where, by definition, $\mathcal{L}_{l,l'}^* = \mathcal{L}_{-l,-l'}$. These spaces can be endowed by the scalar product through the procedure which was explained in the section 3 on the example of Virasoro algebra with central charge $c = -2$. Namely, we

introduce dual field ϕ^* and repeat analysis above for dual representations. Then we identify dual modules of Virasoro algebra with subspaces (or factor spaces) of Fock space by demanding the condition $T^*(\zeta) = T(\zeta)$. Again, choosing the field ϕ as basic field and determining dual field via Riccati type equation, we destroy the symmetry between ϕ and ϕ^* . For instance, Fock submodule $\mathcal{L}_{l,l'}^* = \text{Ker}_{\mathcal{F}_{l,l'}^*}[X^{\star l}]$ will be identified with factor module $\mathcal{L}_{-l,-l'} = \mathcal{F}_{-l,-l'}/\text{Im}_{\mathcal{F}_{l,-l'}}[X^l]$ etc. Thus, we have

Proposition 5.3

Let irreducible representation $\mathcal{L}_{l,l'}$ and dual Verma module $\mathcal{M}_{l,l'}^*$ of Virasoro algebra are realized as bosonic modules $\text{Ker}_{\mathcal{F}_{l,l'}}[X^l]$ and $\mathcal{F}_{-l,-l'}$ correspondingly, where $l = 1, 2, \dots$. Then the scalar product $(\cdot, \cdot)_0$ (3.21) restricted on the vectors from $\mathcal{L}_{l,l'}$ and $\mathcal{M}_{l,l'}^*$ coincides with the following:

$$(\mathbf{u}L_n, \mathbf{v}) = (\mathbf{u}, L_{-n}\mathbf{v}) . \quad (5.10)$$

Let us explain the meaning of this statement. Consider at first vector $\mathbf{u} \in \text{Ker}_{\mathcal{F}_{l,l'}}[X^l] \cong \mathcal{L}_{l,l'}$. Then due to both scalar products $(\cdot, \cdot)_0$ and (5.10) $\mathbf{u}$ has to be orthogonal to any vector $\mathbf{v} \notin \mathcal{F}_{-l,-l'}$. Let now $\mathbf{v} \in \mathcal{F}_{-l,-l'}$. We know from the proposition 5.2 that $\mathcal{F}_{-l,-l'} \cong \mathcal{M}_{l,l'}^*$. It is clear that taking $\mathbf{v} \in \text{Im}_{\mathcal{F}_{l,-l'}}[X^l]$ we would find that this vector is orthogonal to any vector from $\text{Ker}_{\mathcal{F}_{l,l'}}[X^l] \cong \mathcal{L}_{l,l'}$. The reason for this is that subspace $\text{Im}_{\mathcal{F}_{l,-l'}}[X^l]$ is generated by null vector in the $\mathcal{M}_{l,l'}^*$ which is orthogonal to $\mathcal{L}_{l,l'}$. So, the only vectors which have non-trivial scalar product are $\mathbf{u} \in \text{Ker}_{\mathcal{F}_{l,l'}}[X^l]$ and $\mathbf{v} \in \mathcal{F}_{-l,-l'}/\text{Im}_{\mathcal{F}_{l,-l'}}[X^l] \cong \mathcal{L}_{l,l'}^*$. But scalar products (3.21) and (5.10) are the same in these spaces. It should be clear from explicit bosonic realization of generators L_n (5.7).

Now let us turn to consideration of the vertex operator algebra acting in the set of irreducible representations of Vir_c . It is easy to see that such an algebra is generated by the following operators

$$\Phi_{21}^{\pm}(\zeta) : \mathcal{L}_{l,l'} \longrightarrow \mathcal{L}_{l\pm 1,l'} \otimes \mathbf{C}[\zeta] \zeta^{\Delta_{l\pm 1,l'} - \Delta_{l,l'}} , \quad (5.11)$$

$$\Phi_{12}^{\pm}(\zeta) : \mathcal{L}_{l,l'} \longrightarrow \mathcal{L}_{l,l'\pm 1} \otimes \mathbf{C}[\zeta] \zeta^{\Delta_{l,l'\pm 1} - \Delta_{l,l'}} . \quad (5.12)$$

The commutation relations of these operators with the Virasoro generators are given by formulas (3.38) , (3.41) where $\Delta_{1,2}$ and $\Delta_{2,1}$ are defined as in (5.8). In addition, the following conjugation conditions take place

$$\begin{aligned} (\mathbf{u}^* \Phi_{21}^{\star \pm}(\zeta), \mathbf{v}) &= (\mathbf{u}^*, \Phi_{21}^{\mp}(\zeta) \mathbf{v}) , \\ (\mathbf{u}^* \Phi_{12}^{\star \pm}(\zeta), \mathbf{v}) &= (\mathbf{u}^*, \Phi_{12}^{\mp}(\zeta) \mathbf{v}) , \end{aligned} \quad (5.13)$$

where $\mathbf{u}^* \in \mathcal{L}_{l,l'}^*$ and $\mathbf{v} \in \mathcal{L}_{l\pm 1,l'} (\mathcal{L}_{l,l'\pm 1})$.

Now we wish to explain on the example of the operators Φ_{21} how one can find the bosonic realization of such vertex operators. Let us introduce formal operator

$$V_-(\zeta) = \int_C \frac{dz}{2\pi i z} e^{-2i\phi(z)} e^{i\phi(\zeta)}, \quad (5.14)$$

where the integration contour C is chosen as in (3.50). The action of this operator is defined only in the Fock modules $\mathcal{F}_{k,k'}$ with $(\xi + 1)k - \xi k' < 0$ and

$$V_-(\zeta) : \mathcal{F}_{k,k'} \xrightarrow{V_-} \mathcal{F}_{k-1,k'}.$$

The chiral primary operator which increases value k by unit can be obviously constructed as $V_+(\zeta) = e^{i\phi(\zeta)}$:

$$V_+(\zeta) : \mathcal{F}_{k,k'} \xrightarrow{V_+} \mathcal{F}_{k+1,k'}. \quad (5.15)$$

It is easy to see that operators (5.14)-(5.15), satisfy the same commutation relations with Virasoro algebra as Φ_{21} . Moreover, the following proposition holds:

Proposition 5.4

Let l, l' be positive integer numbers such that $(\xi + 1)l - \xi l' < 0$. Then the following diagrams are commutative:

i.

$$\begin{array}{ccc} & \mathcal{F}_{l,l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{l\pm 1,l'} \\ X' l' \downarrow & \downarrow & \downarrow \\ & \mathcal{F}_{l,-l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{l\pm 1,-l'} \end{array} \quad (-1)^{l'} X' l' , \quad (5.16)$$

ii.

$$\begin{array}{ccc} & \mathcal{F}_{-l,-l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{-l\pm 1,-l'} \\ X' l' \uparrow & \uparrow & \uparrow \\ & \mathcal{F}_{-l,l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{-l\pm 1,l'} \end{array} \quad (-1)^{l'} X' l' . \quad (5.17)$$

The propositions 5.1-5.4 ensure that $V_{\pm}$ maps one highest weight representation of Vir_c into another and provide all properties of vertex operators. However, we are capable to identify such bosonic operators with vertex operators Φ_{21} only on the states, where $V_{\pm}$

are defined but not in the whole Fock space. Let $\mathbf{v}$ and $\mathbf{v}^*$ be vectors from $\mathcal{L}_{l,l'}$ and $\mathcal{L}_{l,l'}^*$ correspondingly, then

$$\begin{aligned}
V_+(\zeta)\mathbf{v} &= \Phi_{21}^+(\zeta)\mathbf{v} \quad (l > 0, l' > 0, (\xi+1)l - \xi l' > 0) , \\
V_+(\zeta)\mathbf{v}^* &= \Phi_{21}^{*-}(\zeta)\mathbf{v}^* \quad (l > 1, l' > 0, (\xi+1)l - \xi l' < 0) , \\
V_-(\zeta)\mathbf{v} &= \Phi_{21}^-(\zeta)\mathbf{v} \quad (l > 1, l' > 0, (\xi+1)l - \xi l' < 0) , \\
V_-(\zeta)\mathbf{v}^* &= \Phi_{21}^{*+}(\zeta)\mathbf{v}^* \quad (l > 0, l' > 0, (\xi+1)l - \xi l' > 0) ,
\end{aligned} \tag{5.18}$$

here we explicitly write down in the parenthesis the restrictions on the numbers l, l' of the spaces $\mathcal{L}_{l,l'}$ where the bosonic realization is defined correctly. The bosonization for vertex operators of another type (Φ_{12} and Φ_{12}^*) can be provided at the similar fashion. The result will be the following:

$$\begin{aligned}
V'_+(\zeta)\mathbf{v} &= \Phi_{12}^+(\zeta)\mathbf{v} \quad (l > 0, l' > 0, (\xi+1)l - \xi l' < 0) , \\
V'_+(\zeta)\mathbf{v}^* &= \Phi_{12}^{*-}(\zeta)\mathbf{v}^* \quad (l > 0, l' > 1, (\xi+1)l - \xi l' > 0) , \\
V'_-(\zeta)\mathbf{v} &= \Phi_{12}^-(\zeta)\mathbf{v} \quad (l > 0, l' > 1, (\xi+1)l - \xi l' > 0) , \\
V'_-(\zeta)\mathbf{v}^* &= \Phi_{12}^{*+}(\zeta)\mathbf{v}^* \quad (l > 0, l' > 0, (\xi+1)l - \xi l' < 0) ,
\end{aligned} \tag{5.19}$$

where $\mathbf{v} \in \mathcal{L}_{l,l'}$ and $\mathbf{v}^* \in \mathcal{L}_{l,l'}^*$. The operators $V'_\pm(\zeta)$ reads

$$\begin{aligned}
V'_+(\zeta) &= e^{i\phi'(z)} , \\
V'_-(\zeta) &= \int_C \frac{dz}{2\pi iz} e^{-2i\phi'(z)} e^{i\phi'(\zeta)}
\end{aligned} \tag{5.20}$$

and its properties are quite similar.

The bosonization prescriptions (5.18)-(5.19) together with the conjugation conditions (5.13) allow us to calculate any matrix element of vertex operators Φ_{12} , Φ_{21} . Using the technique described in the previous sections, one can calculate the functions $G_p^\pm$, $G_p'^\pm$ (3.42) for any number ξ and then find the commutation relations in the algebra of vertex operators. We will arrive to the following result [29], [33], [34]:

$$\begin{aligned}
\Phi_{21}^a(\zeta_1)\Phi_{21}^b(\zeta_2)|_{\mathcal{L}_{l,l'}} &= \sum_{c+d=a+b} \mathbf{W} \left[\begin{matrix} l+a+b & l+c \\ l+b & l \end{matrix} \middle| \sigma_{12}, p \right] \Phi_{21}^d(\zeta_2)\Phi_{21}^c(\zeta_1)|_{\mathcal{L}_{l,l'}} , \\
\Phi_{12}^a(\zeta_1)\Phi_{12}^b(\zeta_2)|_{\mathcal{L}_{l,l'}} &= \sum_{c+d=a+b} \mathbf{W} \left[\begin{matrix} l'+a+b & l'+c \\ l'+b & l' \end{matrix} \middle| \sigma_{12}, p' \right] \Phi_{12}^d(\zeta_2)\Phi_{12}^c(\zeta_1)|_{\mathcal{L}_{l,l'}} ,
\end{aligned} \tag{5.21}$$

where $p = e^{i\pi \frac{\xi+1}{\xi}}$ and $p' = e^{i\pi \frac{\xi}{\xi+1}}$. As before, these formulae define the rule of analytical continuation of matrix elements of the vertex operators from the region $|\zeta_1| > |\zeta_2|$ to $|\zeta_2| > |\zeta_1|$ along the contours $C_{\sigma_{12}}$, $\sigma_{12} = \pm$ depicted in fig.2. The nontrivial elements of $\mathbf{W}$ read explicitly:

$$\begin{aligned} \mathbf{W} \left[\begin{array}{cc} l \pm 2 & l \pm 1 \\ l \pm 1 & l \end{array} \middle| \sigma \right] &= p^{\frac{1}{2}\sigma} , \\ \mathbf{W} \left[\begin{array}{cc} l & l \pm 1 \\ l \pm 1 & l \end{array} \middle| \sigma \right] &= \mp \frac{p^{-(\frac{1}{2} \pm l)\sigma}}{[l]_p} , \\ \mathbf{W} \left[\begin{array}{cc} l & l \pm 1 \\ l \mp 1 & l \end{array} \middle| \sigma \right] &= -p^{-\frac{1}{2}\sigma} \frac{[l \pm 1]_p}{[l]_p} , \end{aligned} \quad (5.22)$$

where $[l]_p = \frac{p^l - p^{-l}}{p - p^{-1}}$. It is easy also to find the commutation relation

$$\Phi_{21}^a(\zeta_1) \Phi_{12}^b(\zeta_2) = a \, b \, d(\sigma_{12}) \, \Phi_{12}^b(\zeta_2) \Phi_{21}^a(\zeta_1) , \quad (5.23)$$

with

$$d(\sigma) = e^{-i\frac{\pi}{2}\sigma} . \quad (5.24)$$

Let us discuss the structure of commutation relations. The algebra (5.21) , (5.23) is self-consistent since $d(\sigma_{12}) d(\sigma_{21}) = 1$ ($\sigma_{12} = -\sigma_{21}$) and matrices $\mathbf{W}, \mathbf{W}'$ satisfy to the so-called unitarity condition:

$$\sum_l \mathbf{W} \left[\begin{array}{cc} l_4 & l \\ l_1 & l_2 \end{array} \middle| \sigma_{21} \right] \mathbf{W} \left[\begin{array}{cc} l_4 & l_3 \\ l & l_2 \end{array} \middle| \sigma_{12} \right] = \delta_{l_1, l_3} . \quad (5.25)$$

The associativity condition for this algebra is provided by Yang-Baxter equation in IRF form [33], [34].

$$\begin{aligned} &\sum_l \mathbf{W} \left[\begin{array}{cc} l_4 & l_3 \\ l & l_2 \end{array} \middle| \sigma_{23} \right] \mathbf{W} \left[\begin{array}{cc} l_5 & l_4 \\ l_6 & l \end{array} \middle| \sigma_{13} \right] \mathbf{W} \left[\begin{array}{cc} l_6 & l \\ l_1 & l_2 \end{array} \middle| \sigma_{12} \right] \\ &= \sum_l \mathbf{W} \left[\begin{array}{cc} l_5 & l_4 \\ l & l_3 \end{array} \middle| \sigma_{12} \right] \mathbf{W} \left[\begin{array}{cc} l & l_3 \\ l_1 & l_2 \end{array} \middle| \sigma_{13} \right] \mathbf{W} \left[\begin{array}{cc} l_5 & l \\ l_6 & l_1 \end{array} \middle| \sigma_{23} \right] , \end{aligned} \quad (5.26)$$

and analogously for $\mathbf{W}'$.

3.3 We wish now to construct the representations of ZF algebras associated with algebras of chiral primaries Φ_{21} [8]. It was explained on the simple example $\xi = 1$ (section 3), that to describe representations of ZF algebras one needs involve in the consideration some finite-dimensional algebra U together with Virasoro algebra. Then irreducible representations

of two ZF algebras are decomposed into direct sum of irreducible representations of whole symmetry algebra

$$Symm = Vir_c \otimes U . \quad (5.27)$$

It will be shown that this proposal leads to ZF algebras in general situation ($\xi \neq 1$) too. The finite dimensional subalgebra U of whole symmetry algebra for $\xi \neq 1$ case was considered in the work [29]. It turns to be a direct product of two quantum algebras of the form (2.9). We refer to the notions of quantum algebras and their representations to the section 2.2.

Let us consider the direct product $U_p(sl(2)) \otimes U_{p'}(sl(2))$ of two quantum algebras with generators ($X^\pm, T$) and ($X'^\pm, T'$) correspondingly. The commutations relations of both algebras are determined by (2.9) where the deformation parameters p, p' are given by

$$p = e^{i\pi \frac{\xi+1}{\xi}} , \quad p' = e^{i\pi \frac{\xi}{\xi+1}} . \quad (5.28)$$

We will concentrate on the situation when ξ is a real irrational number more then 1. In this case, as it was noted in section 2, the representations of quantum algebra (2.9) are similar to representations of ordinary $sl(2)$ algebra [26], [27] For this reason, algebra $U_p(sl(2)) \otimes U_{p'}(sl(2))$ admits a set of finite dimensional irreducible representations parameterized by a pair of natural numbers l, l' . We will denote these representations as $\mathcal{V}_{l,l'}$. The basis vectors of space $\mathcal{V}_{l,l'}$ is given by $\mathbf{e}_{l,l'}^{m,m'} = \mathbf{e}_l^m \otimes \mathbf{e}_{l'}^{m'}$, where $\mathbf{e}_l^m$ were described in (2.11).

Let us consider now the whole symmetry algebra $Symm = Vir_c \otimes U_p(sl(2)) \otimes U_{p'}(sl(2))$. The irreducible representation of this algebra is given by tensor product

$$\pi_Z = \oplus_{l,l'} \mathcal{L}_{l,l'} \otimes \mathcal{V}_{l,l'} . \quad (5.29)$$

The scalar product in this space is induced by scalar product in $\mathcal{L}_{l,l'}$ and in $\mathcal{V}_{l,l'}$. We claim that π_Z has the structure of irreducible representation of two ZF algebras. Indeed, let us define the action of generators of ZF algebras on the space (5.29) by the formulae [8] :

$$\begin{aligned} Z_a(\zeta) \mathbf{v} \otimes \mathbf{e}_{l,l'}^{m,m'} &= \\ &= (-1)^{j'-m'} \sum_{b=\pm 1} \begin{pmatrix} \frac{1}{2} & j & j + \frac{b}{2} \\ \frac{a}{2} & m & m + \frac{a}{2} \end{pmatrix}_p \Phi_{21}^b(\zeta) \mathbf{v} \otimes \mathbf{e}_{l+b,l'}^{m+\frac{a}{2},m'} , \end{aligned} \quad (5.30)$$

$$\begin{aligned} Z'_a(\zeta) \mathbf{v} \otimes \mathbf{e}_{l,l'}^{m,m'} &= \\ &= (-1)^{j-m} \sum_{b=\pm 1} \begin{pmatrix} \frac{1}{2} & j' & j' + \frac{b}{2} \\ \frac{a}{2} & m' & m' + \frac{a}{2} \end{pmatrix}_{p'} \Phi_{12}^b(\zeta) \mathbf{v} \otimes \mathbf{e}_{l,l'+b}^{m,m'+\frac{a}{2}} . \end{aligned} \quad (5.31)$$

Here, as usual, $l = 2j + 1, l' = 2j' + 1$. Using precise expressions (2.13) and (5.18)-(5.19) one can find that operators (5.30)-(5.31) satisfy for $\zeta_1 \neq \zeta_2$ the commutation relations:

$$\begin{aligned} Z_a(\zeta_1)Z_b(\zeta_2) &= R_{ab}^{cd}(\sigma_{12}, p) Z_d(\zeta_2)Z_c(\zeta_1) , \\ Z'_a(\zeta_1)Z'_b(\zeta_2) &= R_{ab}^{cd}(\sigma_{12}, p') Z'_d(\zeta_2)Z'_c(\zeta_1) , \\ Z_a(\zeta_1)Z'_b(\zeta_2) &= a b d(\sigma_{12}) Z'_b(\zeta_2)Z_a(\zeta_1) . \end{aligned} \tag{5.32}$$

The meaning of the parameter $\sigma_{12} = \pm$ is the same as in the formula (3.49). The explicit form of the matrix $R_{ab}^{cd}(\sigma_{12}, p)$ is defined by the relations (2.7)-(2.8) .

6. Deformed Virasoro algebra and elliptic ZF algebras.

Now we want to consider a general case of the deformed Virasoro algebra $Vir_{c,x}$ with $0 < x < 1$ and the central charge

$$c = 1 - \frac{6}{\xi(\xi + 1)} , \tag{6.1}$$

where $\xi > 1$ is again an irrational number. Our method, various aspects of which were demonstrated above, is based on the very simple idea. *The structure of representations of deformed Virasoro algebra is determined by the deformed intertwining operators.* The problem of proper deformation of screening operators is not quite evident. Here we will use the deformation suggested in the work [17]. Having the deformed screening operators depending on two continuous parameters x and ξ , we claim that there is a correspondent two parametric algebra $Vir_{c,x}$ which coincides with the Virasoro algebra with central charge (6.1) at the limit $x \rightarrow 1$. Indeed, for given parameters ξ, x any operator constructed from annihilation-creation operators of Fock space belong, by definition, to universal enveloping algebra of deformed Virasoro algebra if it commutes with screening operators. Unfortunately, at present time, we do not know explicit expressions for proper basis of generators of the deformed Virasoro algebra. Therefore the generalization of $\Phi_{21}^{\pm}, \Phi_{12}^{\pm}$ will be based only on properties of the deformed screening operators. In such a way, we obtain the bosonic representation of the elliptic ZF algebra of the deformed vertex operators. We will demonstrate on examples how to provide real computations of matrix elements of the vertex operators. The main point we wish to emphasize in this section is that *the ideas and technique developed in CFT can be generalized for other integrable models.*

6.1. Screening operators for $Vir_{c,x}$ algebra

Let bosonic fields $\phi(z)$ and $\phi'(z)$ are given by the formulas (5.1), (5.3). We will assume that modes b_m satisfy to deformed commutation relations [17]:

$$[b_m, b_n] = m \frac{[m]_x [(\xi + 1)m]_x}{[2m]_x [\xi m]_x} \delta_{m+n,0} , \quad (6.2)$$

and

$$b'_m [m(\xi + 1)]_x = b_m [m\xi]_x . \quad (6.3)$$

Here we use the notation $[a]_x = \frac{x^a - x^{-a}}{x - x^{-1}}$. One can consider these commutation relations as a two-parametric deformation of Heisenberg algebra (3.8) with real parameters $\xi > 1$ and $0 < x < 1$. The meaning of deformation with parameter ξ was discussed in section 5, while the simple example of x-deformation was represented in section 4. The Fock module $\mathcal{F}_p$ for algebra (6.2) can be constructed as an ordinary one. As a vector space it is isomorphic to undeformed Fock space considered in the previous sections because $\mathcal{F}_p$ is still covered by vectors $\oplus_{k=1}^{\infty} \mathbf{C} b_{-n_1} \dots b_{-n_k} \mathbf{f}_p$ with $n_i > 0$. The module $\mathcal{F}_p$ can be endowed by the structure of $\mathbb{Z}$ -graded module if we introduce grading operator as

$$D = \frac{1}{2} \sum_{m>0} \frac{[2m]_x}{[m]_x} (b_{-m} b'_m + b'_{-m} b_m) + \frac{P^2}{2} - \frac{1}{4\xi(\xi + 1)} . \quad (6.4)$$

The x-deformation (6.2) does not change the number of states on every level, and character of the Fock module will be the same as an earlier. Let us note here, that choosing deformation parameter x as real positive number, we are considering x-deformation in generic point. As before the main object of our consideration will be the set of the Fock space $\{\mathcal{F}_{k,k'} \equiv \mathcal{F}_{\frac{(\xi+1)k - \xi k'}{\sqrt{2\xi(\xi+1)}}} \mid k, k' \in \mathbb{Z}\}$. Introduce two-parameters family of formal operators

$$\begin{aligned} X^l &= \int_{C_1} \dots \int_{C_l} \frac{dz_1}{2\pi i z_1} \dots \frac{dz_l}{2\pi i z_l} e^{-i [\phi(z_1 x) + \phi(z_1 x^{-1})]} \dots e^{-i [\phi(z_l x) + \phi(z_l x^{-1})]} , \\ X'^{l'} &= \int_{C_1} \dots \int_{C_{l'}} \frac{dz_1}{2\pi i z_1} \dots \frac{dz_{l'}}{2\pi i z_{l'}} e^{-i [\phi'(z_1 x) + \phi'(z_1 x^{-1})]} \dots e^{-i [\phi'(z_{l'} x) + \phi'(z_{l'} x^{-1})]} . \end{aligned} \quad (6.5)$$

Evidently, these integral operators are deformations of operators (5.5). We have seen example of such deformation (4.1) in section 4. Let us note that X^l and $X'^{l'}$ are related by transformation $\xi \leftrightarrow -1 - \xi$ (or equivalently, $\phi \leftrightarrow \phi'$, $l \leftrightarrow l'$). The worth important fact is that, as in the case $x = 1$, operators X^l and $X'^{l'}$ obey the proposition:

Proposition 6.1 *The action of operator X^l ($X'^{l'}$) is defined only on the Fock modules $\mathcal{F}_{l,k}$ ($\mathcal{F}_{k,l'}$), $k \in \mathbf{Z}$ and $l, l' > 0$. Then*

$$\begin{aligned}
\text{Ker}_{\mathcal{F}_{l,l'}}[X^l] &= \text{Ker}_{\mathcal{F}_{l,l'}}[X'^{l'}] , \\
\text{Ker}_{\mathcal{F}_{l,-l'}}[X^l] &= \text{Ker}_{\mathcal{F}_{l,-l'}}[X'^{l'}] = 0 , \\
\text{Im}_{\mathcal{F}_{l,l'}}[X^l] &= \mathcal{F}_{-l,l'} , \\
\text{Im}_{\mathcal{F}_{l,l'}}[X'^{l'}] &= \mathcal{F}_{l,-l'} , \\
\text{Im}_{\mathcal{F}_{l,-l'}}[X^l] &= \text{Im}_{\mathcal{F}_{-l,l'}}[X'^{l'}] \subset \mathcal{F}_{-l,-l'} .
\end{aligned} \tag{6.6}$$

Now we would like to give the

Definition. *Let operators $P, Q, b_n, n \neq 0$ satisfy the commutation relations (6.2) and screening operators are given by (6.5). The x -deformed Virasoro algebra $\text{Vir}_{c,x}$ is a subalgebra in the universal enveloping algebra of deformed Heisenberg algebra of operators b_n and P . An element from universal enveloping of Heisenberg algebra belongs to deformed Virasoro algebra if it commutes with screening operators.*

The example of the deformed Virasoro algebra was given in section 4. This algebra inherits $\mathbf{Z}$ grading from the algebra (6.2) of operators b_n . Due to definition, screening operators are intertwining operators for representations of deformed Virasoro algebra. Using proposition 6.1 one can investigate the irreducible representations of $\text{Vir}_{c,x}$. Let spaces $\mathcal{L}_{l,l'}$ are determined as in (5.9). We assume that in the generic point of x -deformation the analogue of proposition 5.2 is also correct. It can be rewritten now as:

Conjecture 6.1

Let parameter ξ be irrational number and $\xi > 1$, then

1. *The space $\mathcal{L}_{l,l'}$ for any integer number l, l' is irreducible representation of deformed Virasoro algebra with central charge defined by (3.16). The highest weight vector of $\mathcal{L}_{l,l'}$ coincide with the vector $\mathbf{f}_{l,l'}$ and its weight is given by the formula (5.8).*

2. *For $l \cdot l' \geq 0$ or $l, l' < 0$, Verma modules $\mathcal{M}_{l,l'}$ built upon highest vector $\mathbf{f}_{l,l'}$ coincide with Fock modules $\mathcal{F}_{l,l'}$.*

Now we need make one more assumption concerning to conjugation condition for $\text{Vir}_{c,x}$. Such as in the Fock space there exists a natural scalar product then we expect that the following statement is true:

Conjecture 6.2

Let irreducible representation $\mathcal{L}_{l,l'}$ and dual Verma module $\mathcal{M}_{l,l'}^*$ of deformed Virasoro algebra $\text{Vir}_{c,x}$ are realized as bosonic modules $\text{Ker}_{\mathcal{F}_{l,l'}}[X^l]$ and $\mathcal{F}_{-l,-l'}$ correspondingly,

where $l = 1, 2, \dots$. Then the scalar product $(\cdot, \cdot)_0$ (3.21) restricted on the vectors from $\mathcal{L}_{l,l'}$ and $\mathcal{M}_{l,l'}^*$ confirms with inner conjugation in the deformed Virasoro algebra.

Unfortunately, at present we can not construct proper basis of generators in "quantum" Virasoro algebra and prove assumptions above. Nevertheless, let us try to work out the consequences of these statements. We will see below that such conjugation condition leads to the proper anti-involution in the algebras of chiral primaries.

Let us consider now the generalization of the operators $V_{\pm}(\zeta)$ and $V'_{\pm}(\zeta)$ introduced in the section 5. In the x -deformed case we will define it as follows:

$$\begin{aligned} V_+(\zeta) &= e^{i\phi(\zeta)} , \\ V_-(\zeta) &= \eta^{-1} \int_C \frac{dz}{2\pi i z} e^{-i[(\phi(zx) + \phi(zx^{-1}))]} e^{i\phi(\zeta)} \\ V'_+(\zeta) &= e^{i\phi'(\zeta)} , \\ V'_-(\zeta) &= \eta'^{-1} \int_C \frac{dz}{2\pi i z} e^{-i[(\phi'(zx) + \phi'(zx^{-1}))]} e^{i\phi'(\zeta)} , \end{aligned} \tag{6.7}$$

where the prescription for the integration contours is the same as an earlier. The constants η, η' we will be specified later to provide the convenient normalization of operators (6.7). Note that action of operator V_- (V'_-) is well defined only on the Fock modules $\mathcal{F}_{k,k'}$ with $(\xi + 1)k - \xi k' < 0$ ($(\xi + 1)k - \xi k' > 0$). One can prove the

Proposition 6.2

Let $(\xi + 1)l - \xi l' < 0$ and $l, l' > 0$. Then

- the action of the operators $V_{\pm}(\zeta)$ is described by the commutative diagrams (5.16) (5.17).

- the action of the operators $V'_{\pm}(\zeta)$ is defined by the following commutative diagrams i.

$$\begin{array}{ccc} & \mathcal{F}_{l,l'} & \xrightarrow{V'_{\pm}} \mathcal{F}_{l,l' \pm 1} \\ X^l \downarrow & & \downarrow \\ & \mathcal{F}_{-l,l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{-l,l' \pm 1} \end{array} \quad (-1)^l X^l , \tag{6.8}$$

ii.

$$\begin{array}{ccc} & \mathcal{F}_{-l,-l'} & \xrightarrow{V'_{\pm}} \mathcal{F}_{-l,-l' \pm 1} \\ X^l \uparrow & & \uparrow \\ & \mathcal{F}_{l,-l'} & \xrightarrow{V_{\pm}} \mathcal{F}_{l,-l' \pm 1} \end{array} \quad (-1)^l X^l . \tag{6.9}$$

Let us illustrate this proposition on the simple example.

6.2. Example of calculations

From the proposition 6.1 follows that if Fock module $F_{-1,-1}$ is considered as $Vir_{c,x}$ module then it contains singular vector $X'\mathbf{f}_{-1,1}$. It is easy to show that this vector turns to be proportional to the state $\partial_t \{te^{-i[\phi'(tx)+\phi'(tx^{-1})]}\}|_{t=0}\mathbf{f}_{-1,1} \sim b'_{-1}\mathbf{f}_{-1,-1}$. According to proposition 6.2, operator $V_-(\zeta_1)$ acting on this state set it into null vector in the Verma module $\mathcal{M}_{-2,-1} \cong \mathcal{F}_{-2,-1}$. Due to our conjecture on the scalar product such null vector has to be orthogonal to all states from irreducible module $\mathcal{L}_{2,1}$ and, in particular, to the state $V_+(\zeta_2)\mathbf{f}_{1,1}$:

$$\zeta_1^{-1}f(\zeta_1\zeta_2^{-1}) = (\mathbf{f}_{1,1}V_+(\zeta_2), V_-(\zeta_1)X'\mathbf{f}_{-1,1}) = 0. \quad (6.10)$$

Let us check this formula. As a consequence of bosonic representation it can be represented in the form:

$$\zeta_1^{-1}f(\zeta_1\zeta_2^{-1}) = \partial_t \left\{ t \int_C \frac{dz}{2\pi iz} (\mathbf{f}_{1,1}, e^{i\phi(\zeta_2)} e^{-i\bar{\phi}(z)} e^{i\phi(\zeta_1)} e^{-i\bar{\phi}'(t)} \mathbf{f}_{-1,1})_0 \right\} |_{t=0}, \quad (6.11)$$

where short notations $\bar{\phi}(z)$ and $\bar{\phi}'(z)$ are introduced for $\phi(zx) + \phi(zx^{-1})$ and $\phi'(tx) + \phi'(tx^{-1})$ correspondingly.

The technique of calculation of similar vacuum expectation values in the bosonic space was developed by Dotsenko and Fateev [2]. Let us recall how it works in the case of x-deformed operators. First of all, it is convenient to extract the contribution coming from the zero modes of operators in (6.11). Using the commutation relation for P, Q it is easy to get that ordering of these operators is resulted in the multiple of the form $\zeta_1^{\frac{\xi-1}{4\xi}} \zeta_2^{-\frac{\xi+3}{4\xi}} z^{\frac{1}{\xi}} t^{-1}$. The ordering of oscillator modes b_n , as an ordinary, means that annihilation operators have to stand at right hand side while creation operators should be at left hand side. It is convenient to distinguish here two cases. In the first case one needs provide ordering of oscillators from different exponents while in the second from the same exponent.

First of all, we want to remind the procedure in the first case. Let us elaborate for instance the ordering of the expression $e^{i\phi(\zeta_2)} e^{i\phi(\zeta_1)}$. Using Campbell-Baker-Hausdorff formula $e^A e^B = e^{A+B} e^{\frac{[A,B]}{2}}$ one can find that contributions appearing from coupling of these exponents are equal to

$$g(\zeta_1\zeta_2^{-1}) = e^{-[\phi_+(\zeta_2), \phi_-(\zeta_1)]}, \quad (6.12)$$

where we denote by $\phi_{\pm}$ positive and negative frequency parts of the field ϕ correspondingly. Straightforward computation with using of commutation relations (6.2) leads to the following representation of function $g(\zeta_1\zeta_2^{-1})$:

$$g(z) = \exp \left[- \sum_{m=1}^{+\infty} \frac{[m]_x [m(\xi+1)]_x}{m [2m]_x [m\xi]_x} z^m \right] . \quad (6.13)$$

The sum in this expression converges only for $|z| < 1$. Its analytical continuation on the whole complex plane is given by the following infinite product:

$$g(z) = \frac{(z; x^{2\xi})_{\infty} (x^4 z; x^{2\xi}, x^4)_{\infty} (x^{4+2\xi} z; x^{2\xi}, x^4)_{\infty}}{(x^2 z; x^{2\xi})_{\infty} (x^6 z; x^{2\xi}, x^4)_{\infty} (x^{2+2\xi} z; x^{2\xi}, x^4)_{\infty}} . \quad (6.14)$$

The contributions coming from the averaging of others pairs of exponents can be obtained by the same fashion. Slightly different procedure have to be provided in the ordering of the oscillators in the same exponent. Proceeding as above, one formally get that the ordering of $e^{i\phi(\zeta_1)}$ gives constant ρ such that $\rho^2 = g(1)$. However, as it is seen from (6.14), $g(z)$ has a simple zero at $z = 1$. Let us adopt the following conventions:

$$\rho^2 = \lim_{z \rightarrow 1} \frac{1 - x^2}{1 - z} g(z) . \quad (6.15)$$

Ordinary, one use exponent operators in the normal ordered form from the very beginning. Then there is no need to take care on ordering of oscillators belonging to the same exponent. Making this ordering in the final step is more convenient for us. Indeed, in deformed case this step is resulted in non-trivial constants depending on the deformation parameter that is unlike to the conformal case where renormalization has the similar form for every exponent. We introduce constants like ρ since they provide the natural normalization of operators. Now let us present the final expression for averaging of the integrand in (6.11) :

$$\begin{aligned} (\mathbf{f}_{1,1}, e^{i\phi(\zeta_2)} e^{-i\bar{\phi}(z)} e^{i\phi(\zeta_1)} e^{-i\bar{\phi}'(t)} \mathbf{f}_{-1,1})_0 &= \rho^2 \bar{\rho} \bar{\rho}' \zeta_1^{\frac{\xi-1}{4\xi}} \zeta_2^{-\frac{\xi+3}{4\xi}} z^{\frac{1}{\xi}} t^{-1} \times \\ &g(\zeta_1\zeta_2^{-1}) w(\zeta_1 z^{-1}) w(z\zeta_2^{-1}) u(t\zeta_1^{-1}) u(t\zeta_2^{-1}) \bar{h}(tz^{-1}) . \end{aligned} \quad (6.16)$$

Here $g(z)$ is defined by (6.14), while other functions yield

$$\begin{aligned} w(z) &= \frac{(x^{1+2\xi} z; x^{2\xi})_{\infty}}{(x^{-1} z; x^{2\xi})_{\infty}} , \\ u(z) &= 1 - z , \\ \bar{h}(z) &= \frac{1}{(1 - zx)(1 - zx^{-1})} . \end{aligned} \quad (6.17)$$

As we mentioned above, the constants $\rho, \bar{\rho}, \bar{\rho}'$ are non-trivial functions of deformation parameter x . They are given by:

$$\begin{aligned}\rho^2 &= (1-x^2) \frac{(x^{2\xi}; x^{2\xi})_\infty}{(x^{2+2\xi}; x^{2\xi})_\infty} g^{-1}(x^2) , \\ \bar{\rho}^2 &= (1-x^2) \frac{(x^{-2}; x^{2\xi})_\infty}{(x^{2+2\xi}; x^{2\xi})_\infty} , \\ \bar{\rho}'^2 &= (1-x^2) \frac{(x^2; x^{2+2\xi})_\infty}{(x^{2\xi}; x^{2+2\xi})_\infty} .\end{aligned}\tag{6.18}$$

The integral from expression (6.16) can be expressed in terms of q-special functions. Let us introduce necessary notations. We will need definitions of q-gamma function $\Gamma_q(a)$ and q-hypergeometric function $F_q(a, b, c; z)$. They are usually defined as following

$$\Gamma_q(a) = (1-q)^{1-a} \frac{(q; q)_\infty}{(q^a; q)_\infty} ,\tag{6.19}$$

$$F_q(a, b, c; z) = 1 + \sum_{n=1}^{\infty} \frac{(q^a; q)_n (q^b; q)_n}{(q^c; q)_n (q; q)_n} z^n ,\tag{6.20}$$

here $(q^a; q)_n = \prod_{p=0}^{n-1} (1 - q^{a+p})$. There is the following integral representation for q-hypergeometric function generalizing formula (3.55):

$$\begin{aligned}\int_C \frac{dz}{2\pi i} z^{c-1} \frac{(q^{\frac{1+a}{2}} z^{-1}; q)_\infty}{(q^{\frac{1-a}{2}} z^{-1}; q)_\infty} \frac{(q^{\frac{1+b}{2}} \zeta z; q)_\infty}{(q^{\frac{1-b}{2}} \zeta z; q)_\infty} = \\ = q^{\frac{c(1-a)}{2}} \frac{\Gamma_q(c+a)}{\Gamma_q(c+1) \Gamma_q(a)} F_q(a+c, b, c+1; q^{1-\frac{a+b}{2}} \zeta) .\end{aligned}\tag{6.21}$$

Notice, that in this remarkable formula we have ordinary contour integral rather than Jackson's one. With using of this integral representation the calculation becomes trivial and leads to the formula:

$$\begin{aligned}f(\zeta) = \text{const } g(\zeta) [F_{x^{2\xi}}(2\xi^{-1}, 1 + \xi^{-1}, \xi^{-1}; \zeta x^{-2}) - \\ (1 + \zeta) F_{x^{2\xi}}(1 + 2\xi^{-1}, 1 + \xi^{-1}, 1 + \xi^{-1}; \zeta x^{-2})] .\end{aligned}\tag{6.22}$$

In the special case under the consideration we need the following particular expressions for hypergeometric functions:

$$\begin{aligned}F_q(a, b, b; z) &= \frac{(q^a z; q)_\infty}{(z; q)_\infty} , \\ F_q(2b-2, b, b-1; z) &= (1 + q^{b-1} z) \frac{(q^{2b-1} z; q)_\infty}{(z; q)_\infty} .\end{aligned}\tag{6.23}$$

Substituting these expressions into formula (6.22) one can obtain that $f(\zeta)$ is identical zero. That proves the orthogonality of the vectors $V_-(\zeta_1)X'\mathbf{f}_{-1,1}$ and $V_+(\zeta_2)\mathbf{f}_{1,1}$. Analogously, one can prove that the following matrix element is zero:

$$(\mathbf{f}_{1,1}V'_+(\zeta_2), V'_-(\zeta_1)X\mathbf{f}_{1,-1}) = 0. \quad (6.24)$$

The procedure here is similar to those done before. The only difference is that averaging of the integrand leads to the expression:

$$\begin{aligned} (\mathbf{f}_{1,1}, e^{i\phi'(\zeta_2)} e^{-i\bar{\phi}'(z)} e^{i\phi'(\zeta_1)} e^{-i\bar{\phi}(t)} \mathbf{f}_{1,-1})_0 &= \rho'^2 \bar{\rho}' \bar{\rho} \zeta_1^{\frac{\xi+2}{4(\xi+1)}} \zeta_2^{-\frac{\xi-2}{4(\xi+1)}} z^{-\frac{1}{\xi+1}} t^{-1} \times \\ &g'(\zeta_1 \zeta_2^{-1}) w'(\zeta_1 z^{-1}) w'(z \zeta_2^{-1}) u(t \zeta_1^{-1}) u(t \zeta_2^{-1}) \bar{h}(t z^{-1}). \end{aligned} \quad (6.25)$$

Let us emphasize here that the function $u(\zeta) = e^{[\phi'_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]}$ in this expression is given by the same formula as before, although it is appeared from the ordering of other exponents. This is consequence of the very special form of the proposed x-deformation. Functions $\bar{h}$, $\bar{\rho}$, $\bar{\rho}'^2$ have been written in (6.17) and (6.18) while others have the form

$$\begin{aligned} g'(z) &= \frac{(x^2 z; x^{2+2\xi}, x^4)_\infty (x^{4+2\xi} z; x^{2+2\xi}, x^4)_\infty}{(x^4 z; x^{2+2\xi}, x^4)_\infty (x^{2+2\xi} z; x^{2+2\xi}, x^4)_\infty}, \\ w'(z) &= \frac{(x^{1+2\xi} z; x^{2+2\xi})_\infty}{(xz, x^{2+2\xi})_\infty}, \\ \rho'^2 &= \frac{(x^2; x^{2+2\xi})_\infty}{(x^{2\xi+2}; x^{2+2\xi})_\infty} g'^{-1}(x^2). \end{aligned} \quad (6.26)$$

6.3. Vertex operators for deformed Virasoro algebra

The proposition 6.2 means that operators (6.7) are vertex operators which interpolate between irreducible representations of deformed Virasoro algebra. We assume that bosonic operators $V_\pm$ can be identified with "quantum" analogues of operators $\Phi_{21}^\pm$ by the formulae

$$\begin{aligned} V_+(\zeta)\mathbf{v} &= \Phi_{21}^+(\zeta)\mathbf{v} \quad (l > 0, l' > 0, (\xi+1)l - \xi l' > 0), \\ V_+(\zeta)\mathbf{v}^* &= \Phi_{21}^{*-}(\zeta x^2)\mathbf{v}^* \quad (l > 1, l' > 0, (\xi+1)l - \xi l' < 0), \\ V_-(\zeta)\mathbf{v} &= \Phi_{21}^-(\zeta)\mathbf{v} \quad (l > 1, l' > 0, (\xi+1)l - \xi l' < 0), \\ V_-(\zeta)\mathbf{v}^* &= \Phi_{21}^{*+}(\zeta x^2)\mathbf{v}^* \quad (l > 0, l' > 0, (\xi+1)l - \xi l' > 0). \end{aligned} \quad (6.27)$$

Bosonic realization of operator $\Phi_{12}^\pm$ is determined by similar expressions:

$$\begin{aligned} V'_+(\zeta)\mathbf{v} &= \Phi_{12}^+(\zeta)\mathbf{v} \quad (l > 0, l' > 0, (\xi+1)l - \xi l' < 0), \\ V'_+(\zeta)\mathbf{v}^* &= \Phi_{12}^{*-}(\zeta x^2)\mathbf{v}^* \quad (l > 0, l' > 1, (\xi+1)l - \xi l' > 0), \\ V'_-(\zeta)\mathbf{v} &= \Phi_{12}^-(\zeta)\mathbf{v} \quad (l > 0, l' > 1, (\xi+1)l - \xi l' > 0), \\ V'_-(\zeta)\mathbf{v}^* &= \Phi_{12}^{*+}(\zeta x^2)\mathbf{v}^* \quad (l > 0, l' > 0, (\xi+1)l - \xi l' < 0), \end{aligned} \quad (6.28)$$

If we adjoin the conjugation conditions:

$$\begin{aligned} (\mathbf{v}_1^* \Phi_{21}^{\pm}(\zeta x^2), \mathbf{v}_2) &= (\mathbf{v}_1^*, \Phi_{21}^{\mp}(\zeta) \mathbf{v}_2), \\ (\mathbf{v}_1^* \Phi_{12}^{\pm}(\zeta x^2), \mathbf{v}_2) &= (\mathbf{v}_1^*, \Phi_{12}^{\mp}(\zeta) \mathbf{v}_2), \end{aligned} \quad (6.29)$$

where $\mathbf{v}_1^* \in \mathcal{L}_{l,l'}^*$, $\mathbf{v}_2 \in \mathcal{L}_{l,l'}$ and $|\zeta| = 1$, then formulae (6.27), (6.28) uniquely specify an action of x-deformed vertex operators in irreducible representation of $Vir_{c,x}$. Having the bosonic representation one can get all information on the vertex operators. In principle, our construction is very similar to those known in CFT. The difference appears only in the explicit form of vacuum averaging of exponents. For instance, let us write down the matrix elements of the product of two x-deformed vertex operators. It can be obtained by using the formula (6.21) and knowing coupling of corresponding exponents. If we fix the constants η and η' in definition (6.7) as following:

$$\eta = \eta_{x^{2\xi}}\left(\frac{\xi+1}{\xi}\right), \quad \eta' = \eta_{x^{2+2\xi}}\left(\frac{\xi}{\xi+1}\right), \quad (6.30)$$

where

$$\eta_q^2(a) = (1-x^2) \frac{q^{\frac{a(a-1)}{2}}(1-q)}{\Gamma_q(a)\Gamma_q(1-a)},$$

then we easily find that x-deformation of the functions $G_p^{\pm}$ (3.42) has the form:

$$\begin{aligned} (\mathbf{v}_{l,l'}^*, \Phi_{2,1}^{\pm}(\zeta_2) \Phi_{2,1}^{\mp}(\zeta_1) \mathbf{v}_{l,l'}) &= C \frac{g(\zeta_1 \zeta_2^{-1})}{g(x^2)} G_{x^{2\xi}}\left(\mp \frac{\xi+1}{\xi} l \pm l', \frac{\xi+1}{\xi}; \zeta_1 \zeta_2^{-1}\right), \\ (\mathbf{v}_{l,l'}^*, \Phi_{1,2}^{\pm}(\zeta_2) \Phi_{1,2}^{\mp}(\zeta_1) \mathbf{v}_{l,l'}) &= C' \frac{g'(\zeta_1 \zeta_2^{-1})}{g'(x^2)} G_{x^{2+2\xi}}\left(\mp \frac{\xi}{\xi+1} l' \pm l, \frac{\xi}{\xi+1}; \zeta_1 \zeta_2^{-1}\right). \end{aligned} \quad (6.31)$$

Both formulae here are described by the same function $G_q(c, a; z)$ taken with different parameters. It is given by the expression:

$$G_q(c, a; z) = q^{\frac{(1-a)(2c+a)}{4}} (1-q)^{2a-2} \frac{\Gamma_q(a+c)}{\Gamma_q(c+1)} z^{\frac{a}{4} + \frac{c}{2}} F_q(a+c, a, c+1; q^{1-a} z). \quad (6.32)$$

If we choose the constants η, η' as in (6.30), then the constants C, C' will have the form

$$C = (1-x^2) \Gamma_{x^{2\xi}}\left(\frac{\xi+1}{\xi}\right), \quad C' = \frac{1-x^{2+2\xi}}{\Gamma_{x^{2+2\xi}}\left(\frac{1}{\xi+1}\right)}.$$

This normalization of vertex operators is convenient since it provides the following normalization of functions (6.31) :

$$\begin{aligned} (\mathbf{v}_{l,l'}^*, \Phi_{21}^{\pm}(\zeta_2) \Phi_{21}^{\mp}(\zeta_1) \mathbf{v}_{l,l'}) &= \frac{1-x^2}{1-x^{-2}\zeta_1\zeta_2^{-1}} + \dots, \quad \zeta_1 \rightarrow x^2\zeta_2, \\ (\mathbf{v}_{l,l'}^*, \Phi_{12}^{\pm}(x^2\zeta) \Phi_{12}^{\mp}(\zeta) \mathbf{v}_{l,l'}) &= 1. \end{aligned} \quad (6.33)$$

Further we will see that expressions (6.31) determine the commutation relations of deformed operators likewise to conformal case. For this reason, in order to describe arbitrary matrix element of operators $\Phi_{12}^{\pm}, \Phi_{21}^{\pm}$ it is enough to present the explicit formulae for functions

$$\begin{aligned} \mathfrak{S}_{nm}(\zeta_1, \dots, \zeta_{2n}, \zeta'_1, \dots, \zeta'_{2m}) = & (\mathbf{v}^*_{11}, \Phi_{12}^-(\zeta'_{2m}) \dots \Phi_{12}^-(\zeta'_{m+1}) \Phi_{12}^+(\zeta'_m) \dots \Phi_{1,2}^+(\zeta'_1) \times \\ & \Phi_{21}^-(\zeta_{2n}) \dots \Phi_{21}^-(\zeta_{n+1}) \Phi_{21}^+(\zeta_n) \dots \Phi_{21}^+(\zeta_1) \mathbf{v}_{11}) . \end{aligned} \quad (6.34)$$

To calculate these matrix elements of vertex operators we need know together with the functions (6.14), (6.17), (6.26) also explicit form of the following averaging:

$$\begin{aligned} \bar{g}(\zeta_1 \zeta_2^{-1}) &= e^{-[\bar{\phi}_+(\zeta_2), \bar{\phi}_-(\zeta_1)]} , \\ \bar{g}'(\zeta_1 \zeta_2^{-1}) &= e^{-[\bar{\phi}'_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]} , \\ h(\zeta_1 \zeta_2^{-1}) &= e^{-[\phi_+(\zeta_2), \phi'_-(\zeta_1)]} . \end{aligned} \quad (6.35)$$

Providing standard procedure which was explained above it is not hard to obtain the formulae

$$\begin{aligned} \bar{g}(z) &= (1-z) \frac{(x^{-2}z; x^{2\xi})_{\infty}}{(x^{2+2\xi}z; x^{2\xi})_{\infty}} , \\ \bar{g}'(z) &= (1-z) \frac{(x^2z; x^{2+2\xi})_{\infty}}{(x^{2\xi}z; x^{2+2\xi})_{\infty}} , \\ h(z) &= \frac{(x^3z; x^4)_{\infty}}{(xz; x^4)_{\infty}} . \end{aligned} \quad (6.36)$$

Bosonization technique allows one to represent the functions (6.34) in the form of contour integral from meromorphic functions: ⁴

$$\begin{aligned} \mathfrak{S}_{nm}(\zeta_1, \dots, \zeta_{2n}, \zeta'_1, \dots, \zeta'_{2m}) = & \rho^{2n} (\bar{\rho} \eta^{-1})^n \rho'^{2m} (\bar{\rho}' \eta'^{-1})^m \int_{C_1} \dots \int_{C_n} \prod_k \frac{dz_k}{2\pi i z_k} \times \\ & \int_{S_1} \dots \int_{S_m} \prod_k \frac{dz'_k}{2\pi i z'_k} f_{\frac{\xi+1}{\xi}}^n(\zeta_1, \dots, \zeta_{2n} | z_1, \dots, z_n) f_{\frac{\xi}{\xi+1}}^m(\zeta'_1, \dots, \zeta'_{2m} | z'_1, \dots, z'_m) \\ & \prod_{i < j} g(\zeta_i \zeta_j^{-1}) \prod_{i < j} \bar{g}(z_i z_j^{-1}) \prod_{i < n+j} w(\zeta_i z_j^{-1}) \prod_{n+i \leq j} w(z_i \zeta_j^{-1}) \prod_{i < j} g'(\zeta'_i \zeta'_j{}^{-1}) \prod_{i < j} \bar{g}'(z'_i z'_j{}^{-1}) \times \\ & \prod_{i < m+j} w'(\zeta'_i z'_j{}^{-1}) \prod_{m+i \leq j} w'(z'_i \zeta'_j{}^{-1}) \prod_{i, j} h(\zeta_i \zeta'_j{}^{-1}) \prod_{i, j} \bar{h}(z_i z'_j{}^{-1}) \prod_{i, j} u(\zeta_i z'_j{}^{-1}) \prod_{i, j} u(z_i \zeta'_j{}^{-1}) . \end{aligned} \quad (6.37)$$

⁴ For convenience, we collect all necessary averaging of the exponents in the Appendix.

Here we denoted by symbols $f_{\frac{\xi+1}{\xi}}^n, f_{\frac{\xi}{\xi+1}}^m$ functions representing the contributions of zero modes in averaging of corresponding exponents. Their explicit form is defined by the relations:

$$f_a^n(\zeta_1, \dots, \zeta_{2n} | z_1, \dots, z_n) = \prod_{k=1}^n \zeta_k^{\frac{a}{4}(1+2k)-\frac{1}{2}} \zeta_{n+k}^{\frac{a}{4}(1+2n-2k)-\frac{1}{2}} z_k^{a(k-n-1)+1}. \quad (6.38)$$

The integration contours $\{C_i\}_{i=1}^n$ and $\{S_i\}_{i=1}^m$ are depicted in the fig.6.




Fig. 6: Integration contours.

Note that any contour C_i encloses those singularities of integrand which depend on ζ_k ($2n \geq k \geq n+i$), z_k ($k > i$), ζ'_k ($2n \geq k \geq 1$), z'_k ($n \geq k \geq 1$). At the same time a contour S_i encloses all singularities determining by ζ'_k ($2m \geq k \geq m+i$), z'_k ($k > i$).

6.4. Elliptic ZF algebra of IRF type.

Now let us turn to consideration of commutation relations of vertex operators. They can be derived from the rules of analytical continuation of the functions (6.31) by using the following relations for q-hypergeometric function [35]:

$$\begin{aligned} F_q(a, b, c; z) &= \frac{\Gamma_q(c) \Gamma_q(b-a) \Theta_q(q^a z)}{\Gamma_q(b) \Gamma_q(c-a) \Theta_q(z)} F_q(a, a-c+1, a-b+1; q^{c+1-a-b} z^{-1}) \\ &+ \frac{\Gamma_q(c) \Gamma_q(a-b) \Theta_q(q^b z)}{\Gamma_q(a) \Gamma_q(c-b) \Theta_q(z)} F_q(b, b-c+1, b-a+1; q^{c+1-a-b} z^{-1}) . \end{aligned} \quad (6.39)$$

We have

Proposition 6.3

Vertex operators Φ_{12}, Φ_{21} obey the following commutation relations:

$$\begin{aligned} \Phi_{21}^a(\zeta_1) \Phi_{21}^b(\zeta_2) |_{\mathcal{L}_{l,l'}} &= \sum_{c+d=a+b} \mathbf{W} \left[\begin{matrix} l+a+b & l+c \\ l+b & l \end{matrix} \middle| \frac{\zeta_1}{\zeta_2} \right] \Phi_{21}^d(\zeta_2) \Phi_{21}^c(\zeta_1) |_{\mathcal{L}_{l,l'}} , \\ \Phi_{12}^a(\zeta_1) \Phi_{12}^b(\zeta_2) |_{\mathcal{L}_{l,l'}} &= \sum_{c+d=a+b} \mathbf{W}' \left[\begin{matrix} l'+a+b & l'+c \\ l'+b & l' \end{matrix} \middle| \frac{\zeta_1}{\zeta_2} \right] \Phi_{12}^d(\zeta_2) \Phi_{12}^c(\zeta_1) |_{\mathcal{L}_{l,l'}} , \\ \Phi_{21}^a(\zeta_1) \Phi_{12}^b(\zeta_2) &= a b d \left(\frac{\zeta_1}{\zeta_2} \right) \Phi_{12}^b(\zeta_2) \Phi_{21}^a(\zeta_1) . \end{aligned} \quad (6.40)$$

The function $d(z)$ appeared at the formula above has the form:

$$d(z) = z^{-\frac{1}{2}} \frac{\Theta_{x^4}(xz)}{\Theta_{x^4}(x^3 z)} , \quad (6.41)$$

while matrices $\mathbf{W}, \mathbf{W}'$ can be represented in the form:

$$\begin{aligned} \mathbf{W} \left[\begin{matrix} l_4 & l_3 \\ l_1 & l_2 \end{matrix} \middle| \zeta \right] &= \zeta^{\frac{\xi+1}{2\xi}} \frac{g(\zeta^{-1})}{g(\zeta)} \hat{\mathbf{W}} \left[\begin{matrix} l_4 & l_3 \\ l_1 & l_2 \end{matrix} \middle| \zeta, \frac{\xi+1}{\xi}, x^{2\xi} \right] , \\ \mathbf{W}' \left[\begin{matrix} l'_4 & l'_3 \\ l'_1 & l'_2 \end{matrix} \middle| \zeta \right] &= \zeta^{\frac{\xi}{2(\xi+1)}} \frac{g'(\zeta^{-1})}{g'(\zeta)} \hat{\mathbf{W}} \left[\begin{matrix} l'_4 & l'_3 \\ l'_1 & l'_2 \end{matrix} \middle| \zeta, \frac{\xi}{\xi+1}, x^{2+2\xi} \right] . \end{aligned} \quad (6.42)$$

$\hat{\mathbf{W}}$ denote the following matrices:

$$\begin{aligned} \hat{\mathbf{W}} \left[\begin{matrix} l \pm 2 & l \pm 1 \\ l \pm 1 & l \end{matrix} \middle| \zeta, a, q \right] &= 1 , \\ \hat{\mathbf{W}} \left[\begin{matrix} l & l \pm 1 \\ l \pm 1 & l \end{matrix} \middle| \zeta, a, q \right] &= \zeta^{a(\mp l - 1)} \frac{\Theta_q(q^a) \Theta_q(q^{\mp al} \zeta)}{\Theta_q(q^a \zeta) \Theta_q(q^{\mp al})} , \\ \hat{\mathbf{W}} \left[\begin{matrix} l & l \mp 1 \\ l \pm 1 & l \end{matrix} \middle| \zeta, a, q \right] &= -q^{\mp a^2 l} \zeta^{-a} \frac{\Theta_q(q^{a(\mp l + 1)}) \Theta_q(\zeta)}{\Theta_q(q^{\mp al}) \Theta_q(q^a \zeta)} . \end{aligned} \quad (6.43)$$

The matrices $\mathbf{W}$, $\mathbf{W}'$ satisfy unitarity condition and Yang-Baxter equation in IRF form [3], [36]. These properties provide correspondingly self-consistency and associativity conditions for algebra of vertex operators (6.40). In addition, matrices $\mathbf{W}$ and $\mathbf{W}'$ also satisfy to the crossing symmetry equation:

$$\mathbf{W}' \left[\begin{array}{cc} l'_4 & l'_3 \\ l'_2 & l'_1 \end{array} \middle| x^2 z^{-1} \right] = -\frac{\kappa_{l'_3}}{\kappa_{l'_4}} \mathbf{W}' \left[\begin{array}{cc} l'_2 & l'_4 \\ l'_1 & l'_3 \end{array} \middle| z \right] , \quad (6.44)$$

where

$$\kappa_{l'} = -x^{\frac{\xi(l'-1)}{\xi+1}(\xi l' - 1)} \frac{\Theta_{x^{2+2\xi}}(x^{2\xi l'})}{\Theta_{x^{2+2\xi}}(x^{2\xi})} .$$

Analogous equation holds for the matrix $\mathbf{W}$. Using the commutation relation (6.40) and the property (6.44) one can show that the quadratic combination $\kappa_{l'-1} \Phi_{12}^+(\zeta) \Phi_{12}^-(x^2 \zeta) + \kappa_{l'+1} \Phi_{12}^-(\zeta) \Phi_{12}^+(x^2 \zeta)$ is a central element in the algebra of vertex operators. In the chosen normalization (6.33) we will have:

$$\kappa_{l'-1} \Phi_{12}^+(\zeta) \Phi_{12}^-(x^2 \zeta) + \kappa_{l'+1} \Phi_{12}^-(\zeta) \Phi_{12}^+(x^2 \zeta) = 1 . \quad (6.45)$$

Note that the conjugation (6.29) confirms with the commutation relations (6.40) and condition (6.45). This justifies proposed choice of the scalar product in the conjecture 4.2. and bosonization rules (6.27), (6.28).

6.5. Trigonometric ZF algebra

Let us shortly discuss now limiting cases of x-deformed construction. The first evident limit is conformal one . It can be obtained when parameter x tends to 1 while variable $\ln \zeta$ remains to be finite and non-zero. The bosonization formulae of matrix elements of vertex operators in this limit are obviously equivalent to Dotsenko-Fateev integral representation for conformal blocks in CFT [2]. In particular, one can find that the complicated function $g(z), g'(z), w(z), w'(z)$ at the limit $x \rightarrow 1$, $i \ln z \sim 1$ become correspondingly $(1-z)^{\frac{\xi+1}{2\xi}}$, $(1-z)^{\frac{\xi}{2(\xi+1)}}$, $(1-z)^{\frac{\xi+1}{\xi}}$, $(1-z)^{\frac{\xi}{\xi+1}}$.

Another limiting case corresponds to the case when elliptic matrices $\mathbf{W}$ transforms into trigonometric one [6], [37]. It can be obtained as following. Let us write down the variables x, ζ in the form $z = e^{-i\epsilon\beta}$ and then look at the limit $\epsilon \rightarrow 0$, assuming that β is finite. Notice, that it is convenient to carry out the modular transformation $q = e^{i\pi\tau} \leftrightarrow e^{-i\frac{\pi}{\tau}}$ of theta functions $\Theta_q(\zeta)$ at first. Then the limits can be found straightforwardly. It is not hard to show that functions (6.37), as well as the commutation relations (6.40) are

well-defined in this limit. These expressions can be naturally treated as vacuum averages of certain operators

$$\begin{aligned}\Phi_{21}^{\pm}(\beta) &= \lim_{\epsilon \rightarrow 0} \Phi_{21}^{\pm}(e^{-i\epsilon\beta}) , \\ \Phi_{12}^{\pm}(\alpha) &= \lim_{\epsilon \rightarrow 0} \Phi_{12}^{\pm}(e^{-i\epsilon\alpha}) ,\end{aligned}\tag{6.46}$$

which act in the set of spaces $\lim_{\epsilon \rightarrow 0} \mathcal{L}_{l,l'}$.⁵ Using bosonic realization of the ZF algebra of IRF type, one can construct in this limit representations of ZF algebra of vertex type [23]. Indeed, the finite-dimensional part of total symmetry algebra *Symm* will coincide with those considered in the conformal case, i.e. it is given by the direct product of two quantum groups $U_p(sl(2)) \otimes U_{p'}(sl(2))$. Let us introduce now space π_Z as

$$\pi_Z = \oplus_{l,l'=1}^{\infty} \mathcal{L}_{l,l'} \otimes \mathcal{V}_{l,l'} ,\tag{6.47}$$

where $\mathcal{V}_{l,l'}$ is $l \cdot l'$ -dimensional irreducible representation of the algebra $U_p(sl(2)) \otimes U_{p'}(sl(2))$ and define an action of the operators $Z_{\pm}(\beta)$, $Z'_{\pm}(\alpha)$ by the formula analogous (5.30), (5.31). It is convenient to consider the following simple redefinition of these operators

$$\begin{aligned}Z_{\pm}(\beta) &\rightarrow e^{\mp \frac{\beta}{2\xi}} Z_{\pm}(\beta) , \\ Z'_{\pm}(\alpha) &\rightarrow e^{\mp \frac{\alpha}{2(\xi+1)}} Z'_{\pm}(\alpha) .\end{aligned}\tag{6.48}$$

Then one can show that operators $Z_a(\beta)$, $Z'_a(\alpha)$ generate the ZF algebra of vertex type:

$$\begin{aligned}Z_a(\beta_1)Z_b(\beta_2) &= S_{ab}^{cd}(\beta_1 - \beta_2)Z_d(\beta_2)Z_c(\beta_1), \\ Z'_a(\alpha_1)Z'_b(\alpha_2) &= R_{ab}^{cd}(\alpha_1 - \alpha_2)Z'_d(\alpha_2)Z'_c(\alpha_1), \\ Z_a(\beta)Z'_b(\alpha) &= a \, b \, d(\beta - \alpha)Z'_b(\alpha)Z_a(\beta).\end{aligned}\tag{6.49}$$

The function $d(\beta)$ is the limiting value of the function (6.41). Its explicit form is just $d(\beta) = \tan(\frac{\pi}{4} + i\frac{\beta}{2})$. The matrices $\mathbf{S}_{ab}^{cd}(\beta)$, $\mathbf{R}_{ab}^{cd}(\alpha)$ have the following form:

$$\begin{aligned}\mathbf{S}_{ab}^{cd}(\beta) &= s(\beta) R_{ab}^{cd}(e^{-\frac{2\beta}{\xi}}, e^{i\pi \frac{\xi+1}{\xi}}) , \\ \mathbf{R}_{ab}^{cd}(\alpha) &= r(\alpha) R_{ab}^{cd}(e^{-\frac{2\alpha}{\xi+1}}, e^{i\pi \frac{\xi}{\xi+1}}) ,\end{aligned}\tag{6.50}$$

where nontrivial elements of matrix $R_{ab}^{cd}(t, p)$ given by (2.7). The explicit forms of the functions $r(\alpha)$, $s(\beta)$ is complicate enough and it can be found in the work [17]. ZF algebra (6.49) was introduced in the context of Sin-Gordon model and its physical meaning was discussed in the works [17], [21], [22] .

⁵ To avoid introducing of an additional notation we will denote in this subsection such spaces as $\mathcal{L}_{l,l'}$.

7. Open problems

Now we run into the problem how to construct the representations of the elliptic ZF algebra of vertex type [38]. Let us give an abstract definition of this object. It is a quadratic algebra of the form (6.49) where matrices $\mathbf{R}_{ab}^{cd}, \mathbf{S}_{ab}^{cd}$ have the following form:

$$\begin{aligned} \mathbf{S}_{ab}^{cd}(\beta) &= e^{-i\frac{\xi+1}{2\xi}\epsilon\beta} \frac{g(e^{i\epsilon\beta})}{g(e^{-i\epsilon\beta})} R_{ab}^{cd}(e^{-\frac{2\beta}{\xi}}, e^{i\pi\frac{\xi+1}{\xi}}, e^{-\frac{4\pi}{\epsilon\xi}}), \\ \mathbf{R}_{ab}^{cd}(\alpha) &= e^{-i\frac{\xi}{2(\xi+1)}\epsilon\alpha} \frac{g'(e^{i\epsilon\alpha})}{g'(e^{-i\epsilon\alpha})} R_{ab}^{cd}(e^{-\frac{2\alpha}{\xi+1}}, e^{i\pi\frac{\xi}{\xi+1}}, e^{-\frac{4\pi}{\epsilon(\xi+1)}}). \end{aligned} \quad (7.1)$$

The nontrivial elements of matrix $R_{ab}^{cd}(t, p)$ are defined by the formula (2.2). The function $d(\beta)$ coincides with (6.41) where z is equal to $e^{-i\epsilon\beta}$. Note that matrices (7.1) satisfy the unitarity, crossing symmetry, Yang-Baxter equation and at the limit $\epsilon \rightarrow 0$ they transform to (6.50). As we have studied on the example above, to construct ZF algebra of vertex type from IRF algebra of vertex operators, one need know the comultiplication in the finite-dimensional subalgebra of symmetry algebra. This problem might be very non-trivial since this subalgebra seems to be related with Sklyanin algebra [39]. At the same time, we want to emphasize that the Sklyanin algebra itself is a deformation of $U_p(sl(2))$, while we expect that this new algebra have to be deformation of a tensor product of quantum groups $U_p(sl(2)) \otimes U_{p'}(sl(2))$, to give the limits which are consistent with our constructions in the trigonometric and conformal cases.

8. Acknowledgments

We would like to thank V. Brajnikov for discussion. We are very gratefull to A.B. Zamolodchikov for helpfull comments and his interest in work.

This work was supported by grant DE-FG05-90ER40559. Ya.P. was also supported in part by grant ISF M6N000.

9. Appendix.

In this Appendix we collect the explicit expressions for the functions and constants, which are necessary to compute the matrix elements of vertex operators in formula (6.37).

The functions $g, w, \bar{g}, g', w', \bar{g}', h, u, \bar{h}$ are defined as

$$\begin{aligned}
g(\zeta_1 \zeta_2^{-1}) &= e^{-[\phi_+(\zeta_2), \phi_-(\zeta_1)]} , \\
w(\zeta_1 \zeta_2^{-1}) &= e^{[\phi_+(\zeta_2), \bar{\phi}_-(\zeta_1)]} , \\
\bar{g}(\zeta_1 \zeta_2^{-1}) &= e^{-[\bar{\phi}_+(\zeta_2), \bar{\phi}_-(\zeta_1)]} , \\
g'(\zeta_1 \zeta_2^{-1}) &= e^{-[\phi'_+(\zeta_2), \phi'_-(\zeta_1)]} , \\
w'(\zeta_1 \zeta_2^{-1}) &= e^{[\phi'_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]} , \\
\bar{g}'(\zeta_1 \zeta_2^{-1}) &= e^{-[\bar{\phi}'_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]} , \\
h(\zeta_1 \zeta_2^{-1}) &= e^{-[\phi_+(\zeta_2), \phi'_-(\zeta_1)]} , \\
u(\zeta_1 \zeta_2^{-1}) &= e^{[\phi_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]} , \\
\bar{h}(\zeta_1 \zeta_2^{-1}) &= e^{-[\bar{\phi}_+(\zeta_2), \bar{\phi}'_-(\zeta_1)]} .
\end{aligned} \tag{9.1}$$

They are given by:

$$\begin{aligned}
g(z) &= \frac{(z; x^{2\xi})_\infty (x^4 z; x^{2\xi}, x^4)_\infty (x^{4+2\xi} z; x^{2\xi}, x^4)_\infty}{(x^2 z; x^{2\xi})_\infty (x^6 z; x^{2\xi}, x^4)_\infty (x^{2+2\xi} z; x^{2\xi}, x^4)_\infty} , \\
w(z) &= \frac{(x^{1+2\xi} z; x^{2\xi})_\infty}{(x^{-1} z; x^{2\xi})_\infty} , \\
\bar{g}(z) &= (1 - z) \frac{(x^{-2} z; x^{2\xi})_\infty}{(x^{2+2\xi} z; x^{2\xi})_\infty} , \\
g'(z) &= \frac{(x^2 z; x^{2+2\xi}, x^4)_\infty (x^{4+2\xi} z; x^{2+2\xi}, x^4)_\infty}{(x^4 z; x^{2+2\xi}, x^4)_\infty (x^{2+2\xi} z; x^{2+2\xi}, x^4)_\infty} , \\
w'(z) &= \frac{(x^{1+2\xi} z; x^{2+2\xi})_\infty}{(xz, x^{2+2\xi})_\infty} , \\
\bar{g}'(z) &= (1 - z) \frac{(x^2 z; x^{2+2\xi})_\infty}{(x^{2\xi} z; x^{2+2\xi})_\infty} , \\
h(z) &= \frac{(x^3 z; x^4)_\infty}{(xz; x^4)_\infty} , \\
u(z) &= 1 - z , \\
\bar{h}(z) &= \frac{1}{(1 - zx)(1 - zx^{-1})} .
\end{aligned} \tag{9.2}$$

The constants $\rho, \bar{\rho}, \rho', \bar{\rho}'$ determined as

$$\begin{aligned}
\rho^2 &= \lim_{z \rightarrow 1} \frac{1-x^2}{1-z} g(z) , \\
\bar{\rho}^2 &= \lim_{z \rightarrow 1} \frac{1-x^2}{1-z} \bar{g}(z) , \\
\rho'^2 &= \lim_{z \rightarrow 1} g'(z) , \\
\bar{\rho}'^2 &= \lim_{z \rightarrow 1} \frac{1-x^2}{1-z} \bar{g}'(z) .
\end{aligned} \tag{9.3}$$

have the following values:

$$\begin{aligned}
\rho^2 &= (1-x^2) \frac{(x^{2\xi}; x^{2\xi})_\infty}{(x^{2+2\xi}; x^{2\xi})_\infty} g^{-1}(x^2) , \\
\bar{\rho}^2 &= (1-x^2) \frac{(x^{-2}; x^{2\xi})_\infty}{(x^{2+2\xi}; x^{2\xi})_\infty} , \\
\rho'^2 &= \frac{(x^2; x^{2+2\xi})_\infty}{(x^{2\xi+2}; x^{2+2\xi})_\infty} g'^{-1}(x^2) , \\
\bar{\rho}'^2 &= (1-x^2) \frac{(x^2; x^{2+2\xi})_\infty}{(x^{2\xi}; x^{2+2\xi})_\infty} .
\end{aligned} \tag{9.4}$$

References

- [1] Belavin, A.A., Polyakov, A.M., Zamolodchikov, A.B.: Infinite conformal symmetries in two-dimensional quantum field theory. Nucl. Phys. **B241**, 333-380 (1984)
- [2] Dotsenko, Vl. S. and Fateev, V. A.: Conformal algebra and multipoint correlation functions in 2d statistical models. Nucl. Phys. **B240**[FS12], 312 (1984) ;
Dotsenko, Vl. S. and Fateev, V. A.: Four-point correlation functions and the operator algebra in 2d conformal invariant theories with central charge $c \leq 1$. Nucl. Phys. **B251**[FS13] 691 (1985) .
- [3] Baxter, R.J.: Exactly Solved Models in Statistical Mechanics, Academic Press, London (1982)
- [4] Gervais, J.-L. and Neveu, A.: Novel triangle relation and absense of tachyons in Liouville string theory. Nucl. Phys. B **238**, 125-141 (1984) ;
Gervais, J.-L. and Neveu, A.: New quantum treatment of Liouville field theory B **224**, 329 (1983).
- [5] Tsuchiya, A., Kanie, Y.: Vertex operators in conformal fiels theory on CP^1 and monodromy representations of braid group. Adv.Stud.Pure Math. **16** 297, Kinokunia, Tokyo (1989);
Rehren, K.-H.: Locality of conformal fields in two dimensions: Exchange algebra on the light cone Comm. Math. Phys. **116**, 675-685 (1988);
Froehlich, J.: Statistics of fields, the Yang-Baxter equation, and the theory of Knots and links. In:"Non-perturbative quantum field theory", G. 't Hooft et.al. (eds.), New York: Plenum (1988)
- [6] Zamolodchikov, A.B., Zamolodchikov, Al.B.: Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field theory models. Ann. Phys. (N.Y.) **120**, 253-291 (1979)
- [7] Faddeev, L.D.: Quantum completely integrable models in field theory. Sov. Sci. Rev. Math. Phys. **C1**, 107-155 (1980)
- [8] Moore G., Reshetikhin, N.Yu.: A comment on quantum group symmetry in conformal field theory. Nucl. Phys. **B328**, 557-574 (1989)
- [9] Feigin, B.L., Fuchs, D.B.: Representations of the Virasoro algebra. In: Topology, Proceedings, Leningrad 1982. Faddeev, L.D., Mal'cev, A.A. (eds.). Lecture Notes in Mathematics, vol. **1060**. Berlin Heidelberg, New York: Springer 1984.
- [10] Frenkel, I.B., Reshetikhin, N.Yu.: Quantum affine algebras and holonomic difference equations. Commun. Math. Phys. **146**, 1-60 (1992)
- [11] Foda, O., Miwa, T.: Corner transfer matrices and quantum affine algebras. Int. J. Mod. Phys. **A7**, Suppl. 1A, 279-302 (1992)
- [12] Davies, B., Foda, O., Jimbo, M., Miwa, T., Nakayashiki, A.: Diagonalization of the XXZ Hamiltonian by Vertex Operators. Commun. Math. Phys. **151**, 89-153 (1993)

- [13] Jimbo, M., Miki, K., Miwa, T., Nakayashiki, A.: Correlation functions of the XXZ model for $\Delta < -1$. Phys. Lett. **A168**, 256-263 (1992)
- [14] Foda, O., Jimbo, M., Miwa, T., Miki, K., Nakayashiki, A.: Vertex operators in solvable lattice models. J. math. Phys. **35**, 13-46 (1994)
- [15] Smirnov, F.A.: Form-factors in completely integrable models of quantum field theory. Singapore: World Scientific 1992
- [16] Felder, G.: BRST approach to minimal models. Nucl. Phys. **B317**, 215-236 (1989)
- [17] Lukyanov, S.: Free Field Representation for Massive Integrable Models, Rutgers preprint RU-93-30 (1993) (hep-th/9307196)
- [18] Foda, O., Iohara, K., Jimbo, M., Kedem, R., Miwa, T. and Yan, H.: An elliptic quantum algebra for $\widehat{sl}_2$. Lett. Math. Phys. **32**, 259-268 (1994) ;
Foda, O., Iohara, K., Jimbo, M., Kedem, R., Miwa, T. and Yan, H.: Notes on highest weight modules of the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl}_2)$. (hep-th/9405058)
- [19] Belavin, A.A. : KdV and W . Adv.Stud.Pure Math.**19**, 117 (1989)
- [20] Matsuo, A.: Free Field Representation of Quantum Affine Algebra $U_q(\widehat{sl(2)})$, Nagoya University preprint, August 1992 ;
Shiraishi, J.: Free Boson Representation of $U_q(\widehat{sl(2)})$. UT-617 Tokyo preprint (1992) ;
Kato, A., Quano, Y., Shiraishi, J.: Free Boson Representation of q-Vertex Operators and Their Correlation Functions. UT-618 Tokyo preprint (1992);
Abada, A., Bougourzi, A.H., El Gradechi, M.A.: Deformation of the Wakimoto construction. Mod. Phys. Lett. **A8**, 715-724 (1993)
- [21] Lukyanov, S.: Correlators of the Jost Functions in the Sine-Gordon Model. Phys. Lett. **B325**, 409-417 (1994)
- [22] Lukyanov, S., Shatashvili, S.: Free field representation for the classical limit of the quantum affine algebra. Phys. Lett. **B298**, 111-115 (1993)
- [23] Pasquier, V.: Etiology of IRF models. Comm. Math. Phys. **118**, 355-364 (1988)
- [24] Yang-Baxter equation in integrable systems, ed. Jimbo, M. Singapore: World Scientific (1989)
- [25] Baxter, R.J.: Partition function of the Eight-vertex lattice model. Ann. Phys. **70**, 193-228 (1972)
- [26] Drinfel'd, V.G.: Quantum Groups. In: Proceedings of the International Congress of Mathematics. Berkeley 1986, **1**, pp. 798-820. California: Acad. Press 1987;
Jimbo, M.: A q-difference analogue of $U\mathcal{G}$ and Yang-Baxter equation. Lett. Math. Phys. **10**, 63-68 (1985)
- [27] Kirillov, A.N., Reshetikhin, N.Yu.: Representations of the algebra $U_q(sl(2))$, q-orthogonal polynomials and invariants of links. In: Kac, V.G. (ed.) Infinite-dimensional Lie algebras and groups. Singapore: World Scientific 1989
- [28] Rosso, M.: Finite-dimensional representations of the quantum analog of the enveloping algebra of a complex simple Lie algebra. Comm. Math. Phys. **117**, 581-593 (1988);

- Lusztig, G.: Quantum deformations of certain simple modules over enveloping algebras. *Adv. in Math.* **70**, 237-249 (1988)
- [29] Felder, G., Froehlich, J., Keller, G.: Braid matrices and structure constants for minimal conformal models. *Commun. Math. Phys.* **124**, 647-664 (1989)
- [30] Chaichian, M., Presnajder, P.: Sugawara construction and the q-deformation of Virasoro (super)algebra. *Phys. Lett. B* **277** 109-118 (1992)
- [31] Bernard, D., Felder, J.: Fock representation and BRST cohomology in $SL(2)$ current algebra. *Comm. Math. Phys.* **127**, 145-168 (1990)
- [32] Fateev, V.A., Zamolodchikov, A.B.: Conformal quantum field theory models in two dimensions having $\mathbf{Z}_3$ symmetry. *Nucl.Phys. B* **280** [FS 18], 644-660 (1987) ;
Fateev, V.A., Lukyanov, S.L.: The models of two-dimensional quantum field theory with $\mathbf{Z}_n$ symmetry. *Int.J.Mod.Phys. A* **3**, 507-520 (1988)
- [33] Moore, G., Seiberg, N.: Classical and quantum conformal field theory. *Commun. Math. Phys.* **123**, 177-254 (1989)
- [34] Alvarez-Gaume, L., Gomez, G., Sierra, C.: Quantum group interpretation of some conformal field theories. *Phys. Lett. B* **220**, 142-152 (1989)
- [35] Gasper, G., Rahman, M., Basic hypergeometric series, Cambridge University Press (1990)
- [36] Andrews, G., Baxter, R., Forrester, J.: Eight-vertex SOS model and generalized Rogers-Ramanujan identities. *J. Stat. Phys.* **35**, 193-266 (1984).
- [37] Zamolodchikov, A.B.: Exact two-particle S-matrix of quantum sine-Gordon solitons. *Commun. Math. Phys.* **55**, 183-186 (1977)
- [38] Jimbo, M., Miwa, T., Nakayashiki, A.: Difference equations for the correlation functions of the eight-vertex model. *J. Phys. A: Math. Gen.* **26**, 2199-2209 (1993)
- [39] Sklyanin, E.K.: Some algebraic structures connected with the Yang-Baxter equation. *Funct. Anal. and Appl.* **16** 263-270 (1983);
Sklyanin, E.K.: Some algebraic structures connected with the Yang-Baxter equation. Representations of quantum algebras. *Funct. Anal. and Appl.* **17** 273-284 (1984)